

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 295/08, 223/14, 333/56, 333/54, 307/80, C07C 219/24, 323/30, C07D 307/94, A61K 31/215</p>	<p>A1</p>	<p>(11) International Publication Number: WO 95/08544 (43) International Publication Date: 30 March 1995 (30.03.95)</p>
<p>(21) International Application Number: PCT/CA94/00513 (22) International Filing Date: 23 September 1994 (23.09.94) (30) Priority Data: 08/126,575 24 September 1993 (24.09.93) US (71) Applicant: THE UNIVERSITY OF BRITISH COLUMBIA [CA/CA]; 6328 Memorial Road, Vancouver, British Colum- bia V6T 1Z3 (CA). (72) Inventors: MacLEOD, Bernard, A.; 4036 West 13th Avenue, Vancouver, British Columbia V6R 2T4 (CA). WALKER, Michael, J., A.; 5176 Connaught Drive, Vancouver, British Columbia V6M 3G3 (CA). WALL, Richard, A.; 3181 West 24th Avenue, Vancouver, British Columbia V6L 1R7 (CA). (74) Agents: NASSIF, Omar, A. et al.; McCarthy Tétrault, Suite 4700, Toronto Dominion Bank Tower, Toronto-Dominion Centre, Toronto, Ontario M5K 1E6 (CA).</p>	<p>(81) Designated States: AU, BR, CA, CN, CZ, FI, HU, JP, KR, NO, NZ, PL, RU, UA, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: AMINOCYCLOHEXYLESTERS AND USES THEREOF (57) Abstract Aminocyclohexylester compounds, including thioesters, are disclosed. The compounds of the present invention may be incorporated in compositions and kits. The present invention also discloses a variety of <i>in vitro</i> and <i>in vivo</i> uses for the compounds and compositions, including the treatment of arrhythmias.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

Description

AMINOCYCLOHEXYLESTERS AND USES THEREOF

5

Technical Field

The present invention is generally directed toward aminocyclohexylester compounds, including thioesters, for use in cardiac arrhythmias and the preparation of pharmaceutical compositions and kits.

10

Background of the Invention


Arrhythmia is a variation from the normal rhythm of the heart beat. The major cause of fatalities due to cardiac arrhythmias is the subtype of arrhythmias known as ventricular fibrillation. Conservative estimates indicate that, in the U.S. alone, approximately 300,000 individuals per year suffer heart attacks. Approximately half of these die from sudden cardiac death, the major cause of which is ventricular fibrillation.

Antiarrhythmic agents have been developed to prevent or alleviate cardiac arrhythmia. For example, Class I antiarrhythmic compounds have been used to treat supraventricular arrhythmias and ventricular arrhythmias. Treatment of ventricular arrhythmia is very important since such an arrhythmia, especially ventricular fibrillation, can be fatal. Serious ventricular arrhythmias (ventricular tachycardia and ventricular fibrillation) occur most often in the presence of myocardial ischemia and/or infarction. Ventricular fibrillation often occurs in the setting of acute myocardial ischemia, before infarction fully develops. At present, lidocaine is the current drug of choice for prevention of ventricular fibrillation. However, many Class I antiarrhythmic compounds may actually increase mortality in patients who have had a myocardial infarction. Therefore, there is a need in the art to identify new antiarrhythmic treatments, particularly treatments for ventricular arrhythmias. The present invention fulfills this need, and further provides other related advantages.

Summary of the Invention

Briefly stated, the present invention provides compounds for a variety of uses, including the blockade of ion channels *in vitro* and *in vivo*, and for the treatment of arrhythmias.

In an aspect of the present invention, aminocyclohexyl esters and thioesters are provided. In one embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(I)

wherein X is a direct bond;

or $-(CH_2)_n-Y-$, where $n = 1, 2$, or 3 , and Y is a direct bond, O or S;

10 or $-CH(R_{12})-Y-$, where R_{12} is alkyl of from one to six carbon atoms, a saturated carbocyclic ring of from three to six carbon atoms, phenyl or benzyl, and Y is a direct bond, O or S;


or $-C(R_{13})=CH-$, where R_{13} is hydrogen, alkyl of from one to six carbon atoms, or phenyl;

15

R_1 and R_2 are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

20 or R_1 and R_2 , when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II:



(II)

25


where m is an integer from three to eight, and the ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

5 or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and the heterocyclic ring containing not more than two nitrogen ring atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

10 or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

15 R_3 and R_4 are independently attached to the cyclohexane ring at the 3-, 4-, 5-, or 6-positions, and are independently hydrogen, hydroxy, alkyl of one to six carbon atoms or alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; and


20 A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from formulae III, IV, V, VI, VII or VIII:



(III)


25 where R_5 , R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl, and when X is a direct bond at least one of R_5 , R_6 and R_7 is a hydroxy, fluorine, chlorine, bromine, trifluoromethyl, alkyl of from one to six carbon atoms, or aryl substituent, and when X is $-\text{CH}=\text{CH}-$, and R_1 and R_2 when taken together with the
30 nitrogen atom to which they are attached, form a N-phenylpiperazine ring, and R_3 and R_4 are hydrogen, at least one of R_5 , R_6 and R_7 is a substituent other than hydrogen;

4




(IV)

where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;




(V)

where R₈ and R₉ are defined as above;



(VI)


where R₁₀ is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms;



(VII)

only when X is a direct bond;

5




(VIII)

only when X is a direct bond;

- 5 with the proviso that, when X is $-(CH_2)_n-Y-$, and $n = 1$, and Y is a direct bond, and R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may not be 4-thianaphthenyl.

- 10 In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(I)

15

wherein X is a direct bond;

or $-(CH_2)_n-Y-$, where $n = 1$ and Y is a direct bond, O or S;

or $-CH(R_{12})-$, where R_{12} is alkyl of from one to six carbon atoms;

or $-C(R_{13})=CH-$, where R_{13} is hydrogen;


20

R_1 and R_2 are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

25

or R_1 and R_2 , when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II:



5 where m is an integer from three to eight, and the ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

10 or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and the heterocyclic ring containing not more than two nitrogen ring atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

15 or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

20 R₃ and R₄ are independently attached to the cyclohexane ring at the 3-, 4-, 5-, or 6-positions, and are independently hydrogen, hydroxy, alkyl of one to six carbon atoms or alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; and

25 A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from:

30 formula III where R₅, R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl, and when X is a direct bond at least one of R₅, R₆ and R₇ is a hydroxy, fluorine, chlorine, bromine, trifluoromethyl, alkyl of from one to six carbon atoms, or aryl substituent, and when X is -CH=CH-, and R₁ and R₂, when taken together with the nitrogen atom to which they are attached, form a N-phenylpiperazine

ring, and R_3 and R_4 are hydrogen, at least one of R_5 , R_6 and R_7 is a substituent other than hydrogen;

5 or formula IV where R_8 and R_9 are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;

or formula V where R_8 and R_9 are defined as above;


10 or formula VI where R_{10} is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH_2 , O, S, or $N-R_{11}$ where R_{11} is hydrogen or alkyl of one to six carbon atoms;

15 or formula VII only when X is a direct bond;

or formula VIII only when X is a direct bond;

20 with the proviso that, when X is $-(CH_2)_n-Y-$, and $n = 1$, and Y is a direct bond, and R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may not be 4-thianaphthenyl.

25 In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(I)

30 wherein X is $-(CH_2)_n-Y-$, where $n = 1$ and Y is a direct bond or O; or $-CH(R_{12})-$, where R_{12} is alkyl of from one to six carbon atoms;

R₁ and R₂ are defined as in formula I as first described herein;

R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, alkoxy of one to six carbon atoms, or
5 are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom; and

A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from:

10

formula III where R₅, R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl;

15

or formula IV where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;

20

or formula V where R₈ and R₉ are defined as above;


25

or formula VI where R₁₀ is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms;

30

with the proviso that, when X is -(CH₂)_n-Y-, and n = 1, and Y is a direct bond, and R₁ and R₂, when taken together with the nitrogen atom to which they are attached, form a pyrrolidiny ring, and R₃ and R₄ are hydrogen, A may not be 4-thianaphthenyl.

In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula IX, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(IX)

wherein R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, either form a ring according to formula II, where m is an integer from three to eight, and the ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and the heterocyclic ring containing not more than two nitrogen ring forming atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

R_3 and R_4 are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, methoxy, or are points of attachment of a five-membered oxaspiran ring; and

A is a saturated carbocyclic ring of three to six carbon atoms, or is selected from:

formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R_6 and R_7 is a substituent other than hydrogen;


or formula IV where R_8 and R_9 are hydrogen;

or formula V where R₈ and R₉ are hydrogen;

or formula VI where R₁₀ is hydrogen, and Z is CH₂, O, S, or N-R₁₁ where
 5 R₁₁ is hydrogen or methyl.

In another embodiment, a compound comprises an enantiomer or
 geometric isomer of a compound of formula X, or a solvate or pharmaceutically
 acceptable salt thereof, the compound of the formula:

10



(X)

wherein R₁ and R₂, when taken together with the nitrogen atom to which
 they are attached, form a ring denoted as in formula II, where m is an integer
 15 from three to eight, and the ring may be substituted at any one carbon atom
 by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three
 carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic
 or aliphatic carbocyclic ring of six carbon atoms;

20 or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight
 ring atoms, containing only carbon, nitrogen and optionally oxygen ring
 atoms, and the heterocyclic ring containing not more than two nitrogen ring
 atoms, the second nitrogen being optionally substituted with an alkyl group of
 one to six carbon atoms or a phenyl ring;

25

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl,
 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or
 3-azabicyclo[3.2.0]heptan-3-yl;

R_3 and R_4 are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, methoxy, or are points of attachment of a five-membered oxaspiran ring; and

- 5 A is a saturated carbocyclic ring of from three to six carbon atoms, or is selected from:

- 10 formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R_6 and R_7 is a substituent other than hydrogen;


or formula IV where R_8 and R_9 are hydrogen;

- 15 or formula V where R_8 and R_9 are hydrogen;

or formula VI where R_{10} is hydrogen, and Z is CH_2 , O, S, or N-R_{11} where R_{11} is hydrogen or methyl;

- 20 with the proviso that when R_1 and R_2 , taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may not be 4-thianaphthenyl.

- 25 In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(I)

- 30 wherein X is a direct bond or $-\text{CH}=\text{CH}-$;

R₁ and R₂ are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

- 5 or R₁ and R₂, when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II where m is an integer from three to eight;

- 10 or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

- 15 R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5- positions, and are independently hydrogen, methoxy, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom; and

- 20 A is a saturated carbocyclic ring of from three to six carbon atoms, or is selected from:

formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, trifluoromethyl, methyl or ethyl, and at least one of R₆ and R₇ is a substituent other than hydrogen;

25

or formula IV where R₈ and R₉ are hydrogen;

or formula V where R₈ and R₉ are hydrogen;


- 30 or formula VI where R₁₀ is hydrogen, and Z is O, S, or N-R₁₁ where R₁₁ is hydrogen or methyl;

or formula VII when X is a direct bond;

35

or formula VIII when X is a direct bond.

In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula XI, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(XI)

wherein R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a ring according to formula II where m is an integer from three to eight; and

A is selected from:


formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R_6 and R_7 is a substituent other than hydrogen;

or formula IV where R_8 and R_9 are hydrogen;

or formula V where R_8 and R_9 are hydrogen;

or formula VI where R_{10} is hydrogen, and Z is O or S.

In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula XII, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



wherein R₁ and R₂, when taken together with the nitrogen atom to which they are attached, form a ring according to formula II where m is an integer from three to eight; and

A is selected from:

formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R₆ and R₇ is a substituent other than hydrogen;


or formula IV where R₈ and R₉ are hydrogen;

or formula V where R₈ and R₉ are hydrogen;

or formula VI where R₁₀ is hydrogen, and Z is O or S;

with the proviso that, when R₁ and R₂ taken together with the nitrogen atom to which they are attached form a pyrrolidiny ring, A may not be 4-thianaphthenyl.

In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula XIII, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:




(XIII)

wherein X is a direct bond, *trans*-CH=CH-, -CH₂- or -CH₂-O-;

- 5 R₁ and R₂, when taken together with the nitrogen atom to which they are attached, complete a ring selected from pyrrolidiny, piperidiny, hexahydroazepiny, morpholiny, methylpiperaziny or 3-azabicyclo[3.2.2]nonany; and
- 10 A is selected from cyclohexyl, 3,4-dichlorophenyl, 4-bromophenyl, 1-naphthyl, 2-naphthyl or 3-thianaphthenyl.

In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula XIV, or a solvate or
 15 pharmaceutically acceptable salt thereof, the compound of the formula:



(XIV)


wherein X is a direct bond;

- 20 or -(CH₂)_n-Y-, where n = 1, 2, or 3, and Y is a direct bond, O or S;
 or -CH(R₁₂)-Y-, where R₁₂ is alkyl of from one to six carbon atoms, a saturated carbocyclic ring of from three to six carbon atoms, phenyl or benzyl, and Y is a direct bond, O or S;
- 25 or -C(R₁₃)=CH-, where R₁₃ is hydrogen, alkyl of from one to six carbon atoms, or phenyl;

R₁ and R₂ are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

- 5 or R₁ and R₂, when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II:




- 10 where m is an integer from three to eight, and the ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

- 15 or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and the heterocyclic ring containing not more than two nitrogen ring atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

- 20 or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;


- 25 R₃ and R₄ are independently attached to the cyclohexane ring at the 3-, 4-, 5-, or 6-positions, and are independently hydrogen, hydroxy, alkyl of one to six carbon atoms or alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; and

- 30 A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from formulae III, IV, V, VI, VII or VIII:



(III)


5 where R₅, R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl, and when X is a direct bond at least one of R₅, R₆ and R₇ is a substituent other than hydrogen;



(IV)

10


where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;



(V)

15

where R₈ and R₉ are defined as above;




(VI)

20

where R₁₀ is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms;


25

18



(VII)

only when X is a direct bond;




(VIII)

5

only when X is a direct bond.

10 In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula XIV, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(XIV)

- 15 wherein X is a direct bond;
 or $-(CH_2)_n-Y-$, where $n = 1$, and Y is a direct bond, O or S;
 or $-CH(R_{12})-$, where R_{12} is alkyl of from one to six carbon atoms;
 or $-C(R_{13})=CH-$, where R_{13} is hydrogen;
- 20 R_1 and R_2 are defined as in formula XIV as first described herein;


R_3 and R_4 are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, alkoxy of one to six carbon atoms, or

are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom; and

A is defined as in formula XIV as first described herein.

5

In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula XV, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



10

(XV)

wherein X is a $-\text{CH}_2-$ or $-\text{CH}_2\text{O}-$;

R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a ring according to formula II, where m is an integer from three to eight, and the ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

20

or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and the heterocyclic ring containing not more than two nitrogen ring forming atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

25

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

30

R_3 and R_4 are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, methoxy, or are points of attachment of a five-membered oxaspiran ring; and

- 5 A is a saturated carbocyclic ring of from three to six carbon atoms, or is selected from:

10 formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy;


or formula IV where R_8 and R_9 are hydrogen;

or formula V where R_8 and R_9 are hydrogen;

15

or formula VI where R_{10} is hydrogen, and Z is CH_2 , O, S, or $N-R_{11}$ where R_{11} is hydrogen or methyl.

- 20 In another embodiment, a compound comprises an enantiomer or geometric isomer of a compound of formula XVI, or a solvate or pharmaceutically acceptable salt thereof, the compound of the formula:



(XVI)

- 25 wherein X is a $-CH_2-$ or $-CH_2-O-$;

R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a ring according to formula II where m is an integer from three to eight; and

30

A is selected from:

formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy;

5

or formula IV where R₈ and R₉ are hydrogen;

or formula V where R₈ and R₉ are hydrogen;

10 or formula VI where R₁₀ is hydrogen, and Z is O or S.

In another aspect of the present invention, compositions are provided that comprise an aminocyclohexyl ester or thioester compound described above in combination with a pharmaceutically acceptable carrier or
15 diluent.

The present invention, in another aspect, provides the compounds described above for use in a method for treating arrhythmia in a warm-blooded animal, and for use in methods for blocking ion channels *in vivo* and *in vitro*.


In yet another aspect of the present invention, compounds
20 including those described above are provided for the manufacture of a medicament for the treatment of arrhythmia.

These and other aspects of the present invention will become evident upon reference to the following detailed description.

25 Detailed Description of the Invention

As noted above, the present invention is directed toward aminocyclohexyl ester and thioester compounds which have a variety of uses. Such uses include blockade of ion channels *in vitro* and *in vivo*, and the treatment of arrhythmias.

30 In one aspect, the compounds of the present invention are esters which may be represented by formula I:




(I)

Compounds of formula I are aminocyclohexyl ester compounds. More specifically, these aminocyclohexyl esters are substituted at position 2 of the cyclohexyl ring with an amino group, NR_1R_2 . The cyclohexyl ring may also be substituted with additional substituents (designated as R_3 and R_4) as described in more detail below. Examples of specific embodiments of the elements of compounds represented by formula I include the following.

Depending upon the selection of substituents R_1 and R_2 , the compounds of formula I may be primary, secondary, or tertiary amines (i.e., where R_1 and R_2 both are hydrogen, or only one is hydrogen, or neither are hydrogen, respectively). Where the amine is tertiary, it may be a cyclic amine. Amino substituents R_1 and R_2 may be independently selected from substituents which include hydrogen, alkyl groups containing from three to eight carbon atoms, alkoxyalkyl groups containing from three to eight carbon atoms, and aralkyl groups containing from seven to twelve carbon atoms. As used herein, the term "alkyl group" refers to branched or unbranched saturated hydrocarbon fragments containing the specified number of carbon atoms. Examples include n-propyl, isopropyl, and t-butyl. As used herein, the term "alkoxyalkyl" refers to alkyl groups substituted with alkoxy groups which in turn are alkyl groups attached to an oxygen atom. For example, a methoxyethyl group ($\text{CH}_3\text{OCH}_2\text{CH}_2-$) is a three carbon alkoxyalkyl group. As used herein, the term "aralkyl" refers to alkyl groups substituted with aryl groups. Aryl groups are unsubstituted aromatic groups, with phenyl and naphthyl groups preferred. An example of an aralkyl group is the benzyl group ($\text{C}_6\text{H}_5\text{CH}_2-$) which is a seven carbon aralkyl group.

Alternatively, R_1 and R_2 , when taken together with the amino nitrogen atom, may form a ring represented by formula II:



(II)

where m is an integer from three to eight. Additionally, the ring carbon atoms may be also substituted. Suitable substituents include hydroxy, oxo ($=O$), alkyl groups containing from one to three carbon atoms, and alkoxy groups containing from one to three carbon atoms. The ring may also be fused at two adjacent carbon atoms with a second ring. The second ring may be either an aromatic ring preferably containing either five or six carbon atoms, or an aliphatic carbocycle preferably containing either five or six carbon atoms. Examples of amino substituents containing a fused ring system as described include perhydroindolyl and 1,2,3,4-tetrahydroisoquinoliny groups.

In addition to R_1 and R_2 forming a chain of ring carbon atoms, R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, may complete a saturated monocyclic ring of from five to eight ring atoms which includes additional heteroatoms. Preferably, R_1 and R_2 , when taken together, contain only a single heteroatom. Preferred heteroatoms include nitrogen and oxygen. An example of a ring in which R_1 and R_2 together include oxygen as a ring atom is the morpholiny group. An example of a ring where R_1 and R_2 together include nitrogen as a ring atom is the piperaziny group. For cyclohexyl esters of formula I which are substituted at position 2 with a heterocyclic ring containing two nitrogen ring atoms, the nitrogen of R_1 and R_2 (i.e., the nitrogen ring atom not attached to the cyclohexyl ring) may be further substituted. Suitable substituents include phenyl and alkyl groups containing from one to six carbon atoms. Examples of such N-phenyl or N-alkyl substitution include N-phenylpiperaziny and N-methylpiperaziny, respectively.

Alternatively, R_1 and R_2 , when taken together with the 2-amino nitrogen of formula I, may complete a bicyclic ring. Bicyclic rings include, for example, 3-azabicyclo[3.2.2]nonane, 2-azabicyclo[2.2.2]octane, 3-azabicyclo[3.1.0]hexane, and 3-azabicyclo[3.2.0]heptane. For these derivatives, the 2-substituents of the cyclohexyl esters of formula I are the following groups: 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, and 3-azabicyclo[3.2.0]heptan-3-yl.


Cyclohexane substituents R_3 and R_4 may be independently attached to ring positions 3, 4, 5 or 6 (i.e., both R_3 and R_4 may be attached to the same ring position or each attached to different ring positions). R_3 and R_4 are independently selected from substituents which include hydrogen, hydroxy, alkyl groups containing from one to six carbon atoms, alkoxy groups containing from one to six carbon atoms, or points of attachment of a spiro five- or six-membered heterocyclic ring. Preferred heterocyclic substituents contain either a single oxygen or a single sulfur ring atom.

Depending upon the nature of X, the ester side chain, -X-A in formula I above, may take several forms. A compound of formula I may have X as a direct bond. Alternatively, X may be an alkylene moiety, $-(CH_2)_n-$, where $n = 1, 2$ or 3 , and Y is a direct bond, an oxygen atom, or a sulfur atom; an alkylidene moiety, $-CH(R_{12})-Y-$, where R_{12} is an alkyl group of from one to six carbon atoms, a saturated carbocyclic ring of from three to six carbon atoms (including cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl), a phenyl group, or a benzyl group, and Y is a direct bond, an oxygen atom, or a sulfur atom; or a *cis*- or *trans*-alkenyl moiety, $-C(R_{13})=CH-$, where R_{13} is a hydrogen, an alkyl group of from one to six carbon atoms, or a phenyl group. For compounds of formula I when X is an alkenyl moiety, X is preferably a *trans*-alkenyl moiety.

Ester side chain component A is generally a hydrophobic moiety. Typically, a hydrophobic moiety is comprised of non-polar chemical groups such as hydrocarbons, hydrocarbons substituted with halogens, and ethers. Suitable hydrocarbons include aliphatic and aromatic hydrocarbons. Preferred aliphatic hydrocarbons include alkyl groups containing from five to twelve carbon atoms and carbocyclic groups containing from three to six carbon atoms. Preferred aromatic groups include phenyl, 1-naphthyl, 2-naphthyl, indenyl (including indene where Z, in formula VI below, is CH_2 , indole where Z is $N-R_{11}$, benzofuran where Z is O, and thianaphthene where Z is S), acenaphthyl, and fluorenyl derivatives and are represented by formulae III, IV, V, VI, VII or VIII, respectively. As described below, in a preferred embodiment, Z is O, S or $N-R_{11}$, and in a particularly preferred embodiment Z is O or S.

Phenyl groups within the compounds of the present invention are represented by formula III:


25



(III)

R₅, R₆, and R₇ are independently selected from substituents which include hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl groups containing from one to six carbon atoms, alkoxy groups containing from one to six carbon atoms, or aryl groups. For compounds of formula I when X is a direct bond, at least one of R₅, R₆, and R₇ is a hydroxy, fluorine, chlorine, bromine, trifluoromethyl, alkyl of from one to six carbon atoms, or aryl substituent. For compounds of formula I when X is a -CH=CH-, and R₁ and R₂, when taken together with the nitrogen atom to which they are attached, form a N-phenylpiperazine ring, and R₃ and R₄ are hydrogen, at least one of R₅, R₆ and R₇ is a substituent other than hydrogen.


1-Naphthyl groups within the compounds of the present invention are represented by formula IV:



(IV)

R₈ and R₉ are independently selected from substituents which include hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl groups containing from one to six carbon atoms, alkoxy groups containing from one to six carbon atoms, or aryl groups.


2-Naphthyl groups within the compounds of the present invention are represented by formula V:



(V)

R₈ and R₉ are as defined above.


Indenyl groups of the present invention are represented by formula VI:



(VI)


R₁₀ is selected from substituents which include hydrogen, hydroxy, fluorine,
 5 chlorine, bromine, an alkyl group containing from one to six carbon atoms, an
 alkoxy group containing from one to six carbon atoms, or an aryl group. Z is
 methylene (CH₂), an oxygen atom, sulfur atom, or nitrogen atom bearing a
 substituent R₁₁. The indenyl groups of formula VI are derivatives of indene,
 indole, benzofuran, and thianaphthene when Z is methylene, nitrogen, oxygen,
 10 and sulfur, respectively. Nitrogen substituent R₁₁ may be hydrogen or an alkyl
 group containing from one to six carbon atoms. For compounds of formula I
 when X is -(CH₂)_n-Y-, and n = 1, and Y is a direct bond, and R₁ and R₂, when
 taken together with the nitrogen atom to which they are attached, form a
 pyrrolidiny ring, and R₃ and R₄ are hydrogen, A may not be 4-thianaphthenyl.

15 Acenaphthyl groups within the compounds of the present invention
 are represented by formula VII:



(VII)

20 Fluorenyl groups within the compounds of the present invention
 are represented by formula VIII:




(VIII)

Ester side chain component A may be an acenaphthyl or fluorenyl group only when X is a direct bond. In a preferred embodiment, the acenaphthyl group is a 1-acenaphthyl group, and the fluorenyl group is a 9-fluorenyl group.


As mentioned above, the present invention provides
 5 aminocyclohexyl esters represented by formula I. In a preferred embodiment, where X is $-(CH_2)_n-Y-$, n is 1. For these embodiments, Y is a direct bond, an oxygen atom, or a sulfur atom. In a particularly preferred embodiment, Y is a direct bond or an oxygen atom. In a preferred embodiment, where X is $-CH(R_{12})-$, R_{12} is an alkyl group from one to six carbon atoms. In a preferred
 10 embodiment, where X is $-C(R_{13})=CH-$, R_{13} is a hydrogen atom. For these embodiments, R_3 and R_4 are preferably independently attached to the cyclohexane ring at the 4- or 5- positions.

In a preferred embodiment where X is $-(CH_2)_n-Y-$, where n = 1 and Y is an oxygen atom, and R_1 and R_2 are included in a ring, the present
 15 invention provides aminocyclohexyl esters represented by formula IX:



(IX)

In a preferred embodiment where X is $-(CH_2)_n-Y-$, where n = 1 and Y is a direct bond, and R_1 and R_2 are included in a ring as described above,
 20 the present invention provides aminocyclohexyl esters represented by formula X:




(X)

25 For either formula IX or X, R_3 and R_4 are preferably independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently selected

from substituents which include hydrogen, methoxy, or are points of attachment of a five-membered oxaspiran ring. Similarly, for either formula IX or X, A is preferably a saturated carbocyclic ring of from three to six carbon atoms, or is selected from: formula III where R₅ is hydrogen, and R₆ and R₇ are
5 independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R₆ and R₇ is a substituent other than hydrogen; or formula IV where R₈ and R₉ are hydrogen; or formula V where R₈ and R₉ are hydrogen; or formula VI where R₁₀ is hydrogen, and Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or
10 methyl. However, for compounds represented by formula X, when R₁ and R₂, taken together with the nitrogen atom to which they are attached, form a pyrrolidiny ring, and R₃ and R₄ are hydrogen, A may not be 4-thianaphthenyl.


In a preferred embodiment where X is a direct bond or -CH=CH-, the present invention provides aminocyclohexyl esters represented by formula I.
15 For these embodiments, R₁ and R₂ are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms; or R₁ and R₂, when taken together with the nitrogen atom to which they are attached, form a ring denoted by formula II where m is an integer from three to eight; or R₁ and R₂ complete a ring selected
20 from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl. R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5- positions, and are independently hydrogen, methoxy, or points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom. A is a saturated
25 carbocyclic ring of from three to six carbon atoms or is selected from: formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, trifluoromethyl, methyl, and ethyl, and at least one of R₆ and R₇ is a substituent other than hydrogen; or formula IV where R₈ and R₉ are hydrogen; or formula V where R₈ and R₉ are hydrogen; or formula VI
30 where R₁₀ is hydrogen, and Z is O, S, or N-R₁₁ where R₁₁ is hydrogen or methyl; or formula VII when X is a direct bond; or formula VIII when X is a direct bond.

In a preferred embodiment where X is -(CH₂)_n-Y-, where n = 1 and Y is an oxygen atom, R₁ and R₂ are included in a ring, and there are no R₃
35 and R₄, the present invention provides aminocyclohexyl esters represented by formula XI:



(XI)


- In a preferred embodiment where X is $-(CH_2)_n-Y-$, where $n = 1$ and Y is a direct bond, R_1 and R_2 are included in a ring, and there are no R_3 and R_4 , the present invention provides aminocyclohexyl esters represented by formula XII:



(XII)

- For either formula XI and XII, R_1 and R_2 preferably are taken together with the nitrogen atom to which they are attached to form a ring according to formula II where m is an integer from three to eight. Similarly, for either formula XI or XII, A is preferably selected from: formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R_6 and R_7 is a substituent other than hydrogen; or formula IV where R_8 and R_9 are hydrogen; or formula V where R_8 and R_9 are hydrogen; or formula VI where R_{10} is hydrogen, and Z is O or S.


- In a preferred embodiment where X is selected from a direct bond; *trans*- $C(R_{13})=CH-$ where R_{13} is a hydrogen atom; or $-(CH_2)_n-Y-$ where $n = 1$ and Y is a direct bond or an oxygen atom, R_1 and R_2 are included in a ring, and there are no R_3 and R_4 , the present invention provides aminocyclohexyl esters represented by formula XIII:



(XIII)

R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, complete a ring selected from pyrrolidinyl, piperidinyl, hexahydroazepinyl, morpholinyl, methylpiperazinyl or 3-azabicyclo[3.2.2]nonanyl. A is preferably selected from 3,4-dichlorophenyl, 1-naphthyl, 2-naphthyl, cyclohexyl, 4-bromophenyl, or 3-thianaphthenyl. In a preferred embodiment, the stereochemistry is the trans-configuration. In another preferred embodiment, the stereochemistry is the cis-configuration.

In another aspect, the compounds of the present invention are aminocyclohexyl thioesters which may be represented by formula XIV:




(XIV)

X, R_1 , R_2 , R_3 , R_4 and A are defined as for formula I as first described herein, except that where X is $-(CH_2)_n-Y-$, and $n = 1$, and Y is a direct bond, and R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may be 4-thianaphthenyl.

In a preferred embodiment where X is selected from a direct bond; $-(CH_2)_n-Y-$ where $n = 1$, and Y is a direct bond, an oxygen atom, or a sulfur atom; or $-CH(R_{12})-$ where R_{12} is an alkyl group of from one to six carbon atoms; or $-C(R_{13})=CH-$ where R_{13} is hydrogen; the present invention provides aminocyclohexyl thioesters represented by formula XIV.

R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, alkoxy of from one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom. A is preferably an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of from three to six carbon atoms, or is preferably selected from: formula III where R₅, R₆, and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl, and when X is a direct bond at least one of R₅, R₆ and R₇ is a substituent other than hydrogen; or formula IV where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl; or formula V where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl; or formula VI where R₁₀ is independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl; and Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms; or formula VII when X is a direct bond; a formula VIII when X is a direct bond.

In a preferred embodiment, where X is a $-(CH_2)_n-Y-$, and $n = 1$, and Y is a direct bond or an oxygen atom, and R₁ and R₂ are included in a ring, the present invention provides aminocyclohexyl thioesters represented by formula XV.




(XV)

R₁ and R₂ are preferably taken together with the nitrogen atom to which they are attached to form a ring according to formula II, where m is an integer from three to eight, and the ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic

carbocyclic ring of five to six carbon atoms; or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and the heterocyclic ring containing not more than two nitrogen ring forming atoms, wherein the second nitrogen is
 5 optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring; or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl. R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5- positions, and are independently hydrogen,
 10 methoxy, or are points of attachment of a five-membered oxaspiran ring. A is preferably a saturated carbocyclic ring of from three to six carbon atoms, or is preferably selected from: formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy; or formula IV where R₈ and
 15 R₉ are hydrogen; or formula V where R₈ and R₉ are hydrogen; or formula VI where R₁₀ is hydrogen, and Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or methyl.

In a preferred embodiment where X is -(CH₂)_n-Y-, and n = 1, and Y is a direct bond or an oxygen atom, R₁ and R₂ are included in a ring, and there
 20 are no R₃ and R₄, the present invention provides aminocyclohexyl thioesters represented by formula XVI:



(XVI)

25 R₁ and R₂ are preferably taken together with the nitrogen atom to which they are attached to form a ring according to formula II where m is an integer from three to eight. A is preferably selected from: formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy; or formula IV where
 30 R₈ and R₉ are hydrogen; or formula V where R₈ and R₉ are hydrogen; or formula VI where R₁₀ is hydrogen, and Z is O or S.

In the formulae depicted above, a bond to a substituent and/or a bond that links A with the remainder of a compound may be shown as intersecting one or more bonds in a ring structure. This indicates that the substituent may be attached at any one of the carbon atoms in the ring structure.

5 The wavy line bonds in the above formulae indicate that the substituents at positions 1 and 2 of the cyclohexane ring may be disposed in a *cis* or *trans* relationship. The substituents R₃ and R₄ may also be independently attached *cis* or *trans* to the 2-amino substituent. The compounds of the present invention contain at least two asymmetric carbon atoms and thus exist as
10 enantiomers and diastereomers. Unless otherwise noted, the present invention includes all enantiomeric and diastereomeric forms of the compounds of the above formulae. Pure stereoisomers, mixtures of enantiomers and/or diastereomers, and mixtures of different compounds of the above formulae are included within the present invention.

15 The compounds of the above formulae may be prepared using known synthetic methodology. In general, compounds of the present invention are prepared by reacting the appropriate 1,2-aminocyclohexanol or 1,2-aminocyclohexanethiol with an appropriate carboxylic acid derivative, derived from A-X-COOH. The carboxylic acid may be converted to a reactive
20 intermediate such as the acid chloride by treatment with, for example, thionyl chloride. The reaction between the acid chloride and the alcohol or thiol is carried out in a suitable solvent, such as chloroform. The reaction temperature may be dependent upon the nature of the reactants, however, in general, the reaction may be refluxed. When the reaction has proceeded to substantial
25 completion, the desired product is recovered from the reaction mixture by conventional organic chemistry techniques, and is purified generally by recrystallisation. This reaction sequence as described generates the aminocyclohexylester as the hydrochloride salt, unless an acid scavenger such as triethylamine is added to the mixture, in which case the product is the free base.
30 The hydrochloride may be converted, if desired, to the free base form by known methodologies, and subsequently, if desired, to other acid addition salts by reaction with inorganic or organic acids. Acid addition salts can also be prepared metathetically by reacting one acid addition salt with an acid which is stronger than that of the anion of the initial salt.

35 Alternative routes based on known methodologies, such as those described below, may be used to prepare the aminocyclohexylester compounds of the present invention. The carboxylic acid may be converted to an alternative

reactive derivative such as an activated ester, anhydride or acyl imidazole and reacted with the appropriate 1,2-aminocyclohexanol or 1,2-aminocyclohexanethiol. The carboxylic acid may be reacted directly with the aminoalcohol or aminothiols with the aid of a coupling reagent such as dicyclohexylcarbodiimide. The starting 1,2-aminocyclohexanol or 1,2-aminocyclohexanethiol compounds are prepared by reaction sequences well known in the art. For example, *trans*-1,2-aminocyclohexanols are commonly prepared by the ring-opening reaction of cyclohexene oxide with the appropriate amine. The starting carboxylic acids are known or are prepared by reaction sequences well known in the art.

The synthesis procedures described herein, especially when taken with the general knowledge in the art, provide sufficient guidance to those of ordinary skill in the art to perform the synthesis, isolation, and purification of the preferred compounds described herein and other analogous compounds. Individual enantiomers may be obtained, if desired, from mixtures of the different forms by known methods of resolution, such as the formation of diastereomers, followed by recrystallisation.

The compounds of the above formulae may be in the form of a solvate or a pharmaceutically acceptable salt, e.g., an acid addition salt. Such salts include hydrochloride, sulfate, phosphate, citrate, fumarate, methanesulfonate, acetate, tartrate, maleate, lactate, mandelate, salicylate, succinate and other salts known in the art.

A compound of the present invention may be prepared as a composition by combining it with a pharmaceutically acceptable carrier or diluent. Suitable carriers or diluents include physiological saline. It will be evident to those of ordinary skill in the art that a composition of the present invention may contain more than one aminocyclohexyl ester and/or thioester compound.

As noted above, the present invention provides for utilizing the compounds described above in *in vitro* and *in vivo* methods. In one aspect, ion channels, such as cardiac sodium channels, are blocked *in vitro* or *in vivo*. A preparation that contains ion channels is contacted with, or a warm-blooded animal is administered, an effective amount of an aminocyclohexyl ester and/or thioester compound. Suitable preparations containing cardiac sodium channels include cells isolated from cardiac tissue as well as cultured cell lines. The step of contacting includes, for example, incubation of ion channels with a compound

under conditions and for a time sufficient to permit blockage of the channels by the compound.

In another aspect, the compounds described above are provided for treating arrhythmia. As used herein, "treating arrhythmia" refers to both therapy
5 for arrhythmia and for the prevention of arrhythmias occurring in a heart that is susceptible to arrhythmia. An effective amount of a composition of the present invention is used to treat arrhythmia in a warm-blooded animal, such as a human. Methods of administering effective amounts of antiarrhythmic agents are well known in the art and include the administration of an oral or parenteral dosage
10 form. Such dosage forms include, but are not limited to, parenteral solutions, tablets, capsules, sustained release implants, and transdermal delivery systems. Generally, oral or intravenous administration is preferred. The dosage amount and frequency are selected to create an effective level of the agent without harmful effects. It will generally range from a dosage of from about 0.1 to about
15 100 mg/kg/day, and typically from about 0.1 to 10 mg/kg where administered orally or intravenously for antiarrhythmic effect.

Administration of compositions of the present invention may be carried out in combination with the administration of other agents. For example, it may be desired to administer an opioid antagonist, such as naloxone, if a
20 compound exhibits opioid activity where such activity may not be desired. The naloxone may antagonize opioid activity of the administered compound without adverse interference with the antiarrhythmic activity.

In order to assess whether a compound has a desired pharmacological activity within the present invention, it is subjected to a series
25 of tests. In the first of such tests, a compound is given as increasing (doubling with each dose) intravenous boluses every 8 minutes to a pentobarbital anesthetized rat. The effects of the compound on blood pressure, heart rate and the ECG are measured 30 seconds, 1, 2, 4 and 8 minutes after each dose. Increasing doses are given until the animal dies. The cause of death is identified
30 as being of either respiratory or cardiac origin. This test gives an indication as to whether the compound is blocking sodium channels and/or potassium channels, and in addition gives information about acute toxicity. The indices of sodium channel blockade are increasing P-R interval and QRS widening of the ECG. Potassium channel blockade results in Q-T interval prolongation of the ECG.

35 A second test involves administration of a compound as an infusion to pentobarbital anesthetized rats in which the left ventricle is subjected to electrical square wave stimulation performed according to a preset protocol

described in further detail below. This protocol includes the determination of thresholds for induction of extrasystoles and ventricular fibrillation. In addition, effects on electrical refractoriness are assessed by a single extra beat technique. In addition effects on blood pressure, heart rate and the ECG are recorded. In this test, sodium channel blockers produce the ECG changes expected from the first test. In addition, sodium channel blockers also raise the thresholds for induction of extrasystoles and ventricular fibrillation. Potassium channel blockade is revealed by increasing refractoriness and widening of the Q-T intervals of the ECG.

10 A third test involves exposing isolated rat hearts to increasing concentrations of a compound. Ventricular pressures, heart rate, conduction velocity and ECG are recorded in the isolated heart in the presence of varying concentrations of the compound. This test provides evidence for direct toxic effects on the myocardium. Additionally, selectivity, potency and efficacy of action of a compound can be ascertained under conditions simulating ischaemia. Concentrations found to be effective in this test are expected to be efficacious in the electrophysiological studies.

15 A fourth test is estimation of the antiarrhythmic activity of a compound against the arrhythmias induced by coronary artery occlusion in anaesthetized rats. It is expected that a good antiarrhythmic compound will have antiarrhythmic activity at doses which have minimal effects on either the ECG, blood pressure or heart rate under normal conditions.

20 A compound is also tested directly for effects on sodium and potassium currents in isolated rat myocytes. Isolated rat myocytes are obtained in the conventional manner from isolated hearts. They are used in voltage clamp studies. In order to obtain adequate voltage clamps for estimation of a compound's effects on sodium and potassium currents, the whole-cell patch clamp technique is used. In this technique, a microelectrode is attached to a cell in such a manner that the cell's internal contents are in free communication with the electrode's content. Using the appropriate buffers and conventional voltage step protocols, both sodium and potassium currents can be identified. The activity of a compound is tested on these currents.

25 All of the foregoing tests are performed using rat tissue. In order to ensure that a compound is not having effects which are only specific to rat tissue, further experiments are performed in dogs and primates. In order to assess possible sodium channel and potassium channel blocking actions *in vivo* in dogs, a compound is tested for effects on the ECG, ventricular epicardial

conduction velocity and responses to electrical stimulation. An anesthetized dog is subjected to an open chest procedure to expose the left ventricular epicardium. After the pericardium is removed from the heart a recording/stimulation electrode is sewn onto the epicardial surface of the left ventricle. Using this
5 array, and suitable stimulation protocols, conduction velocity across the epicardium as well as responsiveness to electrical stimulation can be assessed. This information coupled with measurements of the ECG allows one to assess whether sodium and/or potassium channel blockade occurs. As in the first test in rats, a compound is given as a series of increasing bolus doses. At the same time
10 possible toxic effects of a compound on the dog's cardiovascular system is assessed.

The effects of a compound on the ECG and responses to electrical stimulation are also assessed in intact, halothane anesthetized baboons (*Papio anubis*). In this preparation, a blood pressure cannula and ECG electrodes are
15 suitably placed in an anesthetized baboon. In addition, a stimulating electrode is placed into the right ventricle, together with a monophasic action potential electrode. As in the tests described above, ECG and electrical stimulation responses to a compound reveal the possible presence of sodium and/or potassium channel blockade. The monophasic action potential also reveals
20 whether a compound widens the action potential, an action expected of a potassium channel blocker.

The present invention also provides compounds for use in the manufacture of a medicament for the treatment of arrhythmia. Aminocyclohexyl ester and/or thioester compounds including those compounds described above
25 may be used in the manufacture of such a medicament.

The present invention also provides kits that contain a pharmaceutical composition which includes one or more compounds of the above formulae. The kit also includes instructions for the use of the pharmaceutical composition for the blocking of ion channels or for the treatment
30 of arrhythmia. Preferably, a commercial package will contain one or more unit doses of the pharmaceutical composition. For example, such a unit dose may be an amount sufficient for the preparation of an intravenous injection. It will be evident to those of ordinary skill in the art that compounds which are light and/or air sensitive may require special packaging and/or formulation. For example,
35 packaging may be used which is opaque to light, and/or sealed from contact with ambient air, and/or formulated with suitable coatings or excipients.

The following examples are offered by way of illustration and not by way of limitation.

5

EXAMPLES

EXAMPLE 1

(±)-*trans*-[2-(1-pyrrolidiny)cyclohexyl](3,4-dichlorophenoxy)acetate
monohydrochloride
(Compound #1)

(i) Pyrrolidine (10.5 g, 148 mmol), cyclohexene oxide (15 mL, 148 mmol) and water (5 mL) are refluxed under nitrogen until GC or tlc analysis shows the reaction to be complete. The cooled mixture is partitioned between saturated sodium hydroxide solution (150 mL) and ether (150 mL). The aqueous layer is washed with additional ether (75 mL) and the combined ether layers are dried over sodium sulfate. The ether is removed *in vacuo* to leave the crude aminoalcohol (26 g), which is vacuum distilled (bp 51°C at full vacuum) to give (±)-*trans*-[2-(1-pyrrolidiny)]cyclohexanol.

(ii) 3,4-Dichlorophenoxyacetic acid (2.62 g, 12 mmol) is refluxed in thionyl chloride (10 mL) under nitrogen for 1 hour. After stirring at room temperature for a further hour, the thionyl chloride is removed *in vacuo* to leave an orange oil, which is dissolved in chloroform (10 mL). The acid chloride solution is refluxed for 12 hours with a solution of (±)-*trans*-[2-(1-pyrrolidiny)]cyclohexanol (2 g, 12 mmol) in chloroform (5 mL) under nitrogen. Ether (40 mL) is added to the cooled mixture, the mixture is stirred for 30 min., and the crude product is filtered off (3.61 g) and washed with ether. It is recrystallised from the minimum volume of hot methanol/ether to yield the title compound.

EXAMPLE 2

(±)-*trans*-[2-(1-pyrrolidinyl)cyclohexyl](1-naphthoxy)acetate
monohydrochloride
(Compound #2)

5 The title compound is prepared according to the method described in Example 1(ii), using 1-naphthoxyacetic acid (2.4 g, 12 mmol) to prepare the acid chloride as a yellow solid, which is dissolved in chloroform (25 mL). This solution is
10 added to a solution of (±)-*trans*-[2-(1-pyrrolidinyl)]cyclohexanol prepared as in Example 1(i) (2 g, 12 mmol) in chloroform (5 mL) under nitrogen. The mixture is refluxed for 10 hours. All solvent is removed *in vacuo*. The resulting solid is mixed with hot ethyl acetate (100 mL) and methanol is added until it dissolves. On cooling, the solid is collected and washed with ethyl acetate, and
15 recrystallised from hot methanol/ethyl acetate to give the title compound.

EXAMPLE 3

20 (±)-*trans*-{2-[N-(3-azabicyclo[3.2.2]nonyl)]cyclohexyl}(3,4-dichlorophenoxy)acetate monohydrochloride
(Compound #3)

(i) (±)-*trans*-{2-[N-(3-Azabicyclo[3.2.2]nonyl)]}cyclohexanol is prepared
25 according to the method described in Example 1(i) by refluxing 3-azabicyclo[3.2.2]nonane (5 g, 40 mmol), cyclohexene oxide (3.9 mL, 40 mmol) and water (3 mL) for 5 hours. The crude product, a waxy yellowish solid obtained on removal of ether, is used without further purification.

30 (ii) A chloroform solution (15 mL) of 3,4-dichlorophenoxyacetyl chloride, prepared from 3,4-dichlorophenoxyacetic acid (2 g, 9 mmol) as in Example 1(ii), is added to a solution of (±)-*trans*-{2-[N-(3-azabicyclo[3.2.2]nonyl)]}cyclohexanol (2 g, 9 mmol) in chloroform (7 mL) under nitrogen. The mixture is refluxed for 12 hours. Ether (30 mL) is added to
35 the cooled mixture, and the crude product is filtered off and washed with ether. It is recrystallised from the minimum volume of hot ethyl acetate to yield the title compound.

EXAMPLE 4

5 (±)-*trans*-[2-(4-morpholinyl)cyclohexyl](3,4-dichlorophenoxy)acetate
monohydrochloride
(Compound #4)

10 (i) (±)-*trans*-[2-(4-Morpholinyl)]cyclohexanol is prepared by the method
detailed in Example 1(i), refluxing morpholine (5 g, 57 mmol), cyclohexene
oxide (5.8 mL, 57 mmol) and water (3 mL) for 2 hours. The crude product is
purified by full vacuum distillation (bp 75-80°C).

15 (ii) A chloroform solution (10 mL) of 3,4-dichlorophenoxyacetyl chloride,
prepared from 3,4-dichlorophenoxyacetic acid (2.4 g, 11 mmol) as in Example
1(ii), is added to a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol (2 g, 11
mmol) in chloroform (5 mL) under nitrogen. The mixture is refluxed for 8
hours. The crude product is filtered off and washed with ether. More ether may
20 be added to the filtrate to yield further crude product. The product is
recrystallised from hot methanol to give the title compound.

EXAMPLE 5

25 (±)-*trans*-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetate
monohydrochloride
(Compound #5)

30 (i) (±)-*trans*-[2-(1-Hexahydroazepinyl)]cyclohexanol is prepared by the
method detailed in Example 1(i), refluxing hexamethyleneimine (17.3 mL, 0.15
mol), cyclohexene oxide (15.5 mL, 0.15 mol) and water (5 mL). The product is
purified by full vacuum distillation (bp 62-65°C).

35 (ii) A chloroform solution (10 mL) of 3,4-dichlorophenoxyacetyl chloride,
prepared from 3,4-dichlorophenoxyacetic acid (2.36 g, 11 mmol) as in Example
1(ii), is added to a solution of (±)-*trans*-[2-(1-hexahydroazepinyl)]cyclohexanol
(2 g, 10 mmol) in chloroform (5 mL) under nitrogen. The mixture is refluxed for

10 hours. The crude product which precipitates on adding ether to the mixture, is filtered off and washed with ether. It is recrystallised from hot methanol (4 mL), and washed with ether, to give the title compound.

5

EXAMPLE 6

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]benzo[b]thiophene-3-acetate
monohydrochloride

10

(Compound #6)

3-Thianaphtheneacetic acid (4.81 g, 25 mmol) is refluxed in thionyl chloride (16 mL) under nitrogen for 1 hour. After stirring at room temperature for a further 20 min., the thionyl chloride is removed *in vacuo* to leave a brown oil, which is dissolved in chloroform (10 mL). The acid chloride solution is added to a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol prepared as in Example 4(i) (4.5 g, 24.3 mmol) in chloroform (5 mL) under nitrogen. The mixture is refluxed for 11 hours. Ether (25 mL) is added to the cooled mixture and the crude product is filtered off and washed with ether (3 x 10 mL). The product may be recrystallised from hot methanol/ether to give the title compound.

15

20

EXAMPLE 7

25

(±)-*trans*-[2-(1-pyrrolidinyl)cyclohexyl](cyclohexyl)acetate
monohydrochloride
(Compound #7)

The title compound is prepared according to the method described in Example 1(ii), using cyclohexylacetic acid (4.4 g, 31 mmol) to prepare the acid chloride as a pale yellow oil, which is dissolved in chloroform (10 mL). This solution is added to a solution of (±)-*trans*-[2-(1-pyrrolidinyl)]cyclohexanol prepared as in Example 1(i) (5 g, 30 mmol) in chloroform (10 mL) under nitrogen. The mixture is refluxed for 15 hours. The solvent is removed *in vacuo* to leave a solid. The crude product may be recrystallised from hot ethyl acetate or methanol/diethyl ether to give the title compound.

30

35

EXAMPLE 8

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]naphthyl-1-acetate
monohydrochloride
(Compound #8)

1-Naphthylacetic acid (5.30 g, 28.5 mmol) is refluxed in thionyl chloride (10 mL) under nitrogen for 1 hour. After stirring at room temperature for a further 90 min., the thionyl chloride is removed *in vacuo* to leave an orange-brown oil, which is dissolved in chloroform (10 mL). The acid chloride solution is slowly added to a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol prepared as in Example 4(i) (5.0 g, 27 mmol) in chloroform (15 mL) under nitrogen. The mixture is refluxed for 5 hours. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric acid (150 mL) and ether (100 mL). The ether layer is separated and the aqueous is washed with more ether (2 x 50 mL), and then basified by the addition of 50% sodium hydroxide solution. This is then extracted with ether (3 x 50 mL), and the combined ether extracts are washed with water until GC shows no unreacted aminoalcohol in the ether. The ether containing the product is then dried over sodium sulfate, and the solvent is removed to leave the crude free ester. The residue is dissolved in ether (40 mL) and dichloromethane (5 mL) and treated with HCl in ether to precipitate the salt. It is washed with ether and recrystallised from ethyl acetate/methanol to yield the title compound.

EXAMPLE 9

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]naphthyl-2-acetate
monohydrochloride
(Compound #9)

2-Naphthylacetyl chloride is prepared according to the method described in Example 8 using 2-naphthylacetic acid (2.64 g, 14.2 mmol) to give the corresponding acid chloride as a yellow solid which is dissolved in chloroform (10 mL). The solution is refluxed for 9 hours with a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol (2.5 g, 13.5 mmol) in chloroform (10 mL) under

nitrogen. The solvent is removed *in vacuo*, and the residue is partitioned between 1M sodium hydroxide solution (100 mL) and dichloromethane (80 mL). The organic layer is separated and the aqueous is washed with more dichloromethane (2 x 75 mL). The combined dichloromethane extracts are dried
5 over sodium sulfate, and the solvent is removed to leave the crude free ester. This may be purified by column chromatography using silica gel and chloroform/ethyl acetate (3:1) as eluent. The free ester is converted to the salt by treatment with HCl in ether/dichloromethane (3:1). The resulting solid is washed
10 with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 10

15 (±)-*trans*-[2-(4-morpholinyl)cyclohexyl]phenylacetate
 monohydrochloride
 (Compound #10)

Phenylacetyl chloride is prepared according to the method described in Example 8 using phenylacetic acid (2.13 g, 15.6 mmol) to give a pale yellow oil which is
20 dissolved in chloroform (10 mL). This solution is refluxed for 12 hours with a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol (2.75 g, 14.9 mmol) in chloroform (10 mL) under nitrogen. The reaction mixture is partitioned between 1M sodium hydroxide solution (100 mL) and dichloromethane (80 mL). The organic layer is separated and the aqueous is washed with more dichloromethane
25 (2 x 50 mL). The combined dichloromethane extracts are dried over sodium sulfate, and the solvent is removed to leave the crude free ester. This is purified by column chromatography and converted to the hydrochloride salt as described in Example 9. The product is recrystallised from hot methanol to yield the title
30 compound.

EXAMPLE 11

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl](3,4-dichlorophenyl)acetate
monohydrochloride
(Compound #11)

3,4-Dichlorophenylacetyl chloride is prepared according to the method described in Example 8, using 3,4-dichlorophenylacetic acid (2.62 g, 12.8 mmol) to give a yellow oil which is dissolved in chloroform (15 mL). This solution is refluxed for 9 hours with a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol (2.25 g, 12.2 mmol) in chloroform (10 mL) under nitrogen. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric acid (75 mL) and ether (75 mL). The ether layer is separated and the aqueous is washed with more ether (2 x 30 mL), and then basified to pH14 by the addition of 50% sodium hydroxide solution. This is then extracted with ether (4 x 30 mL), and the combined ether extracts are washed with water until GC shows no unreacted aminoalcohol in the ether. The ether containing the product is then dried over sodium sulfate, and the solvent is removed to leave the crude free ester. The residue is dissolved in ether (20 mL) and treated with HCl in ether to precipitate the salt. It is washed with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 12

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl](4-nitrophenyl)acetate
monohydrochloride
(Compound #12)

4-Nitrophenylacetyl chloride, prepared according to the method described in Example 8 using 4-nitrophenylacetic acid (5.4 g, 29.8 mmol), is dissolved in chloroform (15 mL). This solution is refluxed for 9 hours with a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol (5.25 g, 28.3 mmol) in chloroform (15 mL) under nitrogen. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric acid (60 mL) and ether (50 mL). The ether layer is separated and the aqueous layer is washed with more ether (2 x 50 mL), and then basified by the addition of 50% sodium hydroxide solution. This is

then extracted with ether (5 x 60 mL), and the combined ether extracts are washed repeatedly with water to remove any unreacted aminoalcohol. The ether containing the product is then dried over sodium sulfate, and the solvent is removed to leave the crude free ester. The residue is dissolved in ether (20 mL) and treated with HCl in ether to precipitate the salt. It is washed with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 13

(±)-*trans*-[2-(4-methyl-1-piperazinyl)cyclohexyl]naphthyl-2-acetate
dihydrochloride
(Compound #13)

(i) (±)-*trans*-[2-(4-Methyl-1-piperazinyl)]cyclohexanol is prepared by the method detailed in Example 1(i), refluxing 1-methylpiperazine (16.6 mL, 150 mmol), cyclohexene oxide (15.2 mL, 150 mmol) and water (5 mL) for 5 hours. The crude product is purified by full vacuum distillation (bp 94°C) to give a white solid.

(ii) 2-Naphthylacetyl chloride is prepared according to the method described in Example 8 using 2-naphthylacetic acid (2.47 g, 13.3 mmol) to give a yellow solid which is dissolved in chloroform (15 mL). This solution is added to a solution of (±)-*trans*-[2-(4-methyl-1-piperazinyl)]cyclohexanol (2.5 g, 12.6 mmol) in chloroform (10 mL), and the mixture is refluxed for 11 hours under nitrogen. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric acid (100 mL) and ether (60 mL). The ether layer is separated and the aqueous is washed with more ether (2 x 30 mL), and then basified by the addition of 50% sodium hydroxide solution. This is then extracted with ether (4 x 30 mL), and the combined ether extracts are washed repeatedly with water to remove any unreacted aminoalcohol. The ether containing the product is then dried over sodium sulfate, and the solvent is removed to leave the crude free ester. The free ester is dissolved in ether/dichloromethane (35 mL, 4:3) and converted to the dihydrochloride salt by treatment with an excess of HCl in the same solvent. The resulting solid is washed with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 14

5 (±)-*trans*-[2-(4-methyl-1-piperazinyl)cyclohexyl]naphthyl-1-acetate
monohydrochloride
(Compound #14)

1-Naphthylacetyl chloride is prepared according to the method described in
10 Example 8 using 1-naphthylacetic acid (2.47 g, 13.3 mmol) to give a yellow oil
which is dissolved in chloroform (10 mL). This solution is added to a solution of
(±)-*trans*-[2-(4-methyl-1-piperazinyl)]cyclohexanol (2.5 g, 12.6 mmol) in
chloroform (10 mL), and the mixture is refluxed for 11 hours under nitrogen.
The crude free ester is isolated as described in Example 13, and is converted to
15 the monohydrochloride salt in ether/dichloromethane (35 mL, 6:1) by treatment
with 1 equivalent of HCl in the same solvent. The resulting solid is washed with
ether and recrystallised from hot ethyl acetate/methanol to yield the title
compound.

20

EXAMPLE 15

 (±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-1-naphthoate
monohydrochloride
25 (Compound #15)

1-Naphthoyl chloride, prepared according to the method described in Example 8
using 1-naphthoic acid (2.44 g, 14.2 mmol), is dissolved in chloroform (15 mL).
This solution is refluxed for 60 hours with a solution of (±)-*trans*-[2-(4-
30 morpholinyl)]cyclohexanol (2.5 g, 13.5 mmol) in chloroform (12 mL) under
nitrogen. The solvent is removed *in vacuo*, and the residue is treated as in
Example 13 to give the crude free ester. This is purified by column
chromatography using silica gel and chloroform/ethyl acetate (9:1) as eluent.
The free ester is converted to the salt by treatment with HCl in ether. The
35 resulting solid is washed with ether and recrystallised from hot ethyl
acetate/methanol to yield the title compound.

EXAMPLE 16

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]benzo[b]thiophene-4-acetate
monohydrochloride
(Compound #16)

4-Thianaphtheneacetyl chloride is prepared according to the method described in Example 8 using 4-thianaphtheneacetic acid (2.44 g, 12.8 mmol) to give a red-brown oil which is dissolved in chloroform (15 mL). This solution is refluxed for 27 hours with a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol (2.27 g, 13.5 mmol) in chloroform (10 mL) under nitrogen. The solvent is removed *in vacuo*, and the residue is treated as in Example 13 to give the crude free ester. This is purified by column chromatography using silica gel and chloroform/ethyl acetate (9:1) as eluent. The free ester is converted to the salt by treatment with HCl in ether. The resulting solid is washed with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 17

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl](4-bromophenyl)acetate
monohydrochloride
(Compound #17)

4-Bromophenylacetyl chloride is prepared according to the method described in Example 8 using 4-bromophenylacetic acid (2.75 g, 12.8 mmol) to give an oil which is dissolved in chloroform (10 mL). This solution is refluxed for 19 hours with a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol (2.25 g, 12.2 mmol) in chloroform (10 mL) under nitrogen. The solvent is removed *in vacuo*, and the residue is treated as in Example 13 to give the crude free ester, which is then dissolved in 1M hydrochloric acid (70 mL). Sodium chloride (14 g) is added to the solution which is then extracted with chloroform (60 mL then 2 x 40 mL). The combined extracts are dried over sodium sulfate and the solvent removed *in vacuo*. The residue is recrystallised from hot ethyl acetate/methanol to give the title compound.

EXAMPLE 18

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-3,4-dichlorocinnamate
monohydrochloride
(Compound #18)

3,4-Dichlorocinnamyl chloride is prepared according to the method described in Example 8 using 3,4-dichlorocinnamic acid (2.46 g, 11.4 mmol) to give an oil which is dissolved in chloroform (10 mL). This solution is refluxed for 35 hours with a solution of (\pm)-*trans*-[2-(4-morpholinyl)]cyclohexanol (2.0 g, 10.8 mmol) in chloroform (10 mL) under nitrogen. The solvent is removed *in vacuo* and the residue partitioned between 10% aqueous sodium hydroxide (50 mL) and dichloromethane (50 mL). The aqueous layer is extracted with more dichloromethane (2 x 50 mL) and the combined extracts are dried over sodium sulfate. The solvent is removed and the crude ester is purified by column chromatography using silica gel and chloroform/ethyl acetate (1:1) as eluent. The free ester is converted to the salt by treatment with HCl in dichloromethane/ether (1:4). The resulting solid is washed with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 19

25 **(±)-*trans*-[2-(1-piperidinyl)cyclohexyl]thiophenoxyacetate
monohydrochloride
(Compound #19)**

30 (i) (\pm) -*trans*-[2-(1-Piperidinyl)]cyclohexanol is prepared by the method detailed in Example 1(i), refluxing piperidine (14.8 mL, 0.15 mol), cyclohexene oxide (15.5 mL, 0.15 mol) and water (5 mL). The product is purified by full vacuum distillation to give a colorless liquid.

(ii) A chloroform (10 mL) solution of thiophenoxyacetyl chloride, prepared from thiophenoxyacetic acid (1.85 g, 11 mmol) as in Example 8, is refluxed for 12 hours with a solution of (±)-*trans*-[2-(1-piperidiny)]cyclohexanol (1.85 g, 10 mmol) in chloroform (15 mL) under nitrogen. The crude product, which

precipitates on cooling the mixture, is filtered off and washed with ether. It is recrystallised from hot methanol (4 mL), and washed with ether, to give the title compound.

5

EXAMPLE 20

(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-2-(1-naphthyl)propionate
monohydrochloride
(Compound #20)

10

(i) 1-Naphthylacetic acid (10 g, 53.7 mmol) is dissolved in dry THF (150 mL) under nitrogen. The solution is cooled to -10°C, treated with 1.6M butyl lithium in hexanes (72 mL, 115 mmol), and stirred for 15 minutes. A solution of
15 iodomethane (3.7 mL, 59.4 mmol) in dry THF (50 mL) is added, and the mixture is allowed to warm to room temperature and stir for several hours. The solvent is removed *in vacuo*, and the residue dissolved in 5% sodium hydroxide solution (500 mL), washed with ether (3 x 200 mL), and then acidified by the addition of
20 6N hydrochloric acid. The aqueous is then extracted with dichloromethane (3 x 200 mL), the organic extracts are dried over sodium sulfate, and the solvent removed to leave 2-(1-naphthyl)propionic acid as a white solid.

(ii) 2-(1-Naphthyl)propionyl chloride, prepared according to the method described in Example 8 using 2-(1-naphthyl)propionic acid (2.55 g, 12.8 mmol), is
25 dissolved in chloroform (10 mL). The acid chloride solution is added to a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol prepared as in Example 4(i) (2.25 g, 12.2 mmol) in chloroform (10 mL) under nitrogen. The mixture is refluxed for 14 hours. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric acid (150 mL) and ether (100 mL). The
30 ether layer is separated and the aqueous is washed with more ether (2 x 50 mL), and then basified by the addition of 50% sodium hydroxide solution. This is then extracted with ether (3 x 50 mL), and the combined ether extracts are washed with water until GC shows no unreacted aminoalcohol in the ether. The
35 ether containing the product is then dried over sodium sulfate, and the solvent is removed to leave the crude free ester. The residue is dissolved in ether (40 mL) and dichloromethane (5 mL) and treated with HCl in ether to precipitate the salt.

It is washed with ether and recrystallised from ethyl acetate/methanol to yield the title compound.

5

EXAMPLE 21

(±)-(1 α ,2 β ,4 β ,5 β)-[4,5-dimethoxy-2-(1-pyrrolidiny)cyclohexyl]benzofuran-2-acetate monohydrochloride
(Compound #21)

10

(i) Iodine (122 g, 480 mmol) is added gradually over 5 hours to a mixture of 1,4-cyclohexadiene (77 g, 963 mmol) and sodium iodate (51.5 g, 240 mmol) in glacial acetic acid (50 mL). This mixture is stirred for a further 1 hour and then water (17.3 g, 963 mmol) and potassium acetate (94.3 g, 963 mmol) are added and the mixture is refluxed for 3 hours. The acetic acid is removed by vacuum distillation, and the residue is dissolved in ether (400 mL) and washed with saturated sodium metabisulfite solution (400 mL). The aqueous layer is extracted with ethyl acetate (3 x 80 mL), and the combined organic extracts are dried over sodium sulfate and the solvent removed. The resulting oil is dissolved in ether (200 mL) and treated with 10% potassium hydroxide in methanol until chromatography shows the conversion to *cis*-4,5-cyclohexenediol to be complete. The mixture is neutralized by the addition of solid carbon dioxide. The solvent is removed and the product is partitioned between saturated sodium chloride solution and ethyl acetate. The ethyl acetate extracts are dried over sodium sulfate, the solvent is removed and the product, *cis*-4,5-cyclohexenediol, is purified by vacuum distillation (bp 90-100°C at full vacuum) to give a white solid.

(ii) A solution of the diol (37.1 g, 325 mmol) in dry THF (600 mL) is added over 1 hour to a cooled, stirred suspension of sodium hydride (32 g of a 60% dispersion in mineral oil, 800 mmol) in dry THF (800 mL) under nitrogen. A solution of methyl iodide (60 mL, 964 mmol) in THF (60 mL) is added and the mixture stirred for 12 hours. The reaction is quenched by the addition of moist ether followed by water. The solvent is removed and the residue partitioned between water (200 mL) and ether (300 mL). The aqueous layer is extracted with additional ether (2 x 75 mL) and the combined organic extracts are dried over sodium sulfate and the solvent removed. The product, 4,5-

dimethoxycyclohexene, is purified by vacuum distillation (bp 90°C at 20 mmHg) to give a colorless oil.

(iii) m-Chloroperbenzoic acid (43 g, 250 mmol) is dissolved in ether (400 mL), any aqueous layer is separated, and the ether solution is dried over sodium sulfate and added quickly dropwise to a solution of 4,5-dimethoxycyclohexene (17.7 g, 125 mmol) in ether (250 mL). The mixture is stirred for 12 hours, most of the ether is removed (450 mL) *in vacuo*, and dichloromethane is added (400 mL). The solution is washed with 20% sodium metabisulfite solution (3 x 200 mL), saturated sodium bicarbonate solution (3 x 200 mL) and brine (3 x 200 mL). The solvent is removed and the epoxide is refluxed with pyrrolidine (40 mL, 480 mmol) and water (10 mL) for 2 hours. After cooling, 25% sodium hydroxide solution (30 mL) is added and the excess pyrrolidine removed by distillation. The product, 4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexanol, is extracted into dichloromethane (3 x 100 mL) and purified by distillation (bp 90-100°C at full vacuum).

(iv) Benzofuran-2-acetyl chloride, prepared according to the method described in Example 8 using benzofuran-2-acetic acid (2.1 g, 12 mmol), is dissolved in chloroform (15 mL). This solution is added to a solution of 4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexanol (2.5 g, 11 mmol) in chloroform (10 mL), and the mixture is refluxed for 11 hours under nitrogen. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric acid (100 mL) and ether (60 mL). The ether layer is separated and the aqueous is washed with more ether (2 x 30 mL), and then basified by the addition of 50% sodium hydroxide solution. This is then extracted with ether (4 x 30 mL), and the combined ether extracts are washed repeatedly with water to remove any unreacted aminoalcohol. The ether containing the product is then dried over sodium sulfate, and the solvent is removed to leave the crude free ester. The free ester is converted to the hydrochloride salt by treatment with HCl in ether. The resulting solid is washed with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 22

(±)-*trans*-{2-[bis(2-methoxyethyl)amino]cyclohexyl}fluorene-9-carboxylate
monohydrochloride
(Compound #22)

- 5
- (i) The intermediate aminoalcohol, (±)-*trans*-{2-[bis(2-methoxyethyl)amino]}cyclohexanol, is prepared by the method detailed in Example 1 (i), refluxing bis(2-methoxyethyl)amine (25 mL, 169 mol),
10 cyclohexene oxide (17.2 mL, 170 mmol) and water (8 mL) for 30 hours. The crude product is purified by full vacuum distillation (bp 83-85°C) to give a clear liquid.
- (ii) 9-Fluorenicarboxylic acid (2.43 g, 11.5 mmol) is converted to the
15 corresponding acid chloride according to the method described in Example 8. The resulting white solid is dissolved in chloroform (10 mL) and refluxed under nitrogen with a chloroform solution (10 mL) of (±)-*trans*-{2-[bis(2-methoxyethyl)amino]}cyclohexanol (2.7 g, 11 mmol) for 20 hours. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric
20 acid (100 mL) and ether (100 mL). The ether layer is separated and the aqueous is washed with more ether (2 x 50 mL), and then basified by the addition of 50% sodium hydroxide solution. This is then extracted with ether (4 x 40 mL), and the combined ether extracts are washed repeatedly with water to remove any unreacted aminoalcohol. The ether containing the product is then dried over
25 sodium sulfate, and the solvent is removed to leave the crude free ester. The free ester is converted to the hydrochloride salt by treatment with HCl in ether. The resulting solid is washed with ether and recrystallised from hot methanol to yield the title compound.

EXAMPLE 23

S-{{(\pm)-*trans*-[2-(4-morpholinyl)cyclohexyl]}(3,4-dimethoxyphenyl)thioacetate
monohydrochloride
(Compound #23)

- 5
- (i) Morpholine (7.6 g, 87 mmol), cyclohexene sulfide (10 g, 88 mmol) and water (5 mL) are refluxed under nitrogen for 5 hours or until GC or tlc analysis show the reaction to be complete. The cooled mixture is partitioned between saturated sodium hydroxide solution (150 mL) and dichloromethane (150 mL). The aqueous layer is washed with dichloromethane (2 x 50 mL) and the combined organic layers are dried over sodium sulfate. The solvent is removed *in vacuo* to give (\pm)-*trans*-[2-(4-morpholinyl)]cyclohexanethiol which is purified by full vacuum distillation (bp 90-100°C).
- 10
- (ii) (3,4-dimethoxyphenyl)acetyl chloride is prepared according to the method described in Example 8 using 3,4-dimethoxyphenylacetic acid (2.85 g, 14.5 mmol) to give a brown oil which is dissolved in chloroform (10 mL). This solution is refluxed for 3 hours with a solution of (\pm)-*trans*-[2-(4-morpholinyl)]cyclohexanethiol (2.8 g, 14 mmol) in chloroform (10 mL) under nitrogen. The reaction mixture is partitioned between 1M sodium hydroxide solution (100 mL) and dichloromethane (80 mL). The organic layer is separated and the aqueous is washed with more dichloromethane (2 x 50 mL). The combined dichloromethane extracts are dried over sodium sulfate, and the solvent is removed to leave the crude free ester. This is purified by column chromatography and converted to the hydrochloride salt by treatment with HCl in ether. The product is recrystallised from hot ethyl acetate/methanol to yield the title compound.
- 15
- 20
- 25
- 30
- Compounds prepared according to the methods described above were generally characterized by proton and/or carbon-13 NMR and/or infrared spectroscopy. Elemental analysis was obtained for compounds listed in Table 1.

EXAMPLE 24

S- $\{(\pm)$ -*trans*-[2-(4-morpholinyl)cyclohexyl]}naphthyl-1-thioacetate
monohydrochloride
(Compound #24)

1-Naphthylacetyl chloride is prepared according to the method described in Example 8 using 1-naphthylacetic acid (2.0 g, 10.8 mmol) to give a yellow oil which is dissolved in chloroform (10 mL). This solution is added to a solution of
10 $\{(\pm)$ -*trans*-[2-(4-morpholinyl)]cyclohexanethiol (2 g, 10 mmol) in chloroform (10 mL), and the mixture is stirred at room temperature for 8 hours under nitrogen. The solvent is removed and the residue is partitioned between 1M hydrochloric acid (100 mL) and ether (80 mL). The organic layer is separated and the aqueous is washed with more ether (2 x 60 mL). The aqueous mixture is
15 basified to pH>12 by the addition of 50% sodium hydroxide solution, and extracted with ether (1 x 80 mL, 2 x 50 mL). These ether extracts of the basic aqueous solution are washed repeatedly with water to remove any unreacted aminothiol, dried over sodium sulfate, and the solvent is removed to leave the crude free thioester. This is converted to the hydrochloride salt by dissolving it
20 in 1M hydrochloric acid (60 mL), adding sodium chloride (12 g), and extracting with chloroform (3 x 75 mL). The chloroform is removed *in vacuo* and the residue is recrystallised from ethyl acetate/methanol to yield the title compound.

25

EXAMPLE 25

S- $\{(\pm)$ -*trans*-{2-[bis(2-methoxyethyl)amino]cyclohexyl}}-4-bromothiobenzoate
monohydrochloride
(Compound #25)

30

(i) The intermediate aminothiol, $\{(\pm)$ -*trans*-{2-[bis(2-methoxyethyl)amino]}cyclohexanethiol, is prepared by the method detailed in Example 23(i), refluxing bis(2-methoxyethyl)amine (13 mL, 88 mmol), cyclohexene sulfide (10 g, 88 mmol) and water (8 mL).

35

(ii) 4-Bromobenzoyl chloride (2.4 g, 11 mmol) in chloroform (10 mL) is added to a solution of $\{(\pm)$ -*trans*-{2-[bis(2-

methoxyethylamino]]cyclohexanethiol (2 g, 10 mmol) in chloroform (10 mL), and the mixture is stirred at room temperature for 8 hours under nitrogen. The crude free thioester is obtained using the work-up procedure described in Example 24, and is converted to the hydrochloride salt by dissolving it in 1M hydrochloric acid (60 mL), adding sodium chloride (12 g), and extracting with chloroform (3 x 75 mL). The chloroform is removed *in vacuo* and the residue is recrystallised from ethyl acetate/methanol to yield the title compound.

10

EXAMPLE 26

(±)-*trans*-[2-(diisopropylamino)cyclohexyl]acenaphthene-1-carboxylate
monohydrochloride
(Compound #26)

15

(i) The intermediate aminoalcohol, (±)-*trans*-(2-diisopropylamino)cyclohexanol, is prepared by the method detailed in Example 1(i), refluxing diisopropylamine (22.3 mL, 170 mmol), cyclohexene oxide (17.2 mL, 170 mmol) and water (8 mL) for 30 hours. The crude product is purified by full vacuum distillation.

20

(ii) Acenaphthene-1-carboxylic acid (2.28 g, 11.5 mmol) is prepared according to the method described by P.R. Halfpenny et al. in *Synthesis*, 1990, 517-519 or in U.S. Patent Nos. 4,906,655 and 5,019,588. It is converted to the corresponding acid chloride according to the method described in Example 8, and is dissolved in chloroform (10 mL) and refluxed under nitrogen with a solution of (±)-*trans*-(2-diisopropylamino)cyclohexanol (2.2 g, 11 mmol) for 20 hours. The solvent is removed *in vacuo*, and the residue is partitioned between 1M hydrochloric acid (100 mL) and ether (100 mL). The ether layer is separated and the aqueous is washed with more ether (2 x 50 mL), and then basified by the addition of 50% sodium hydroxide solution. This is then extracted with ether (4 x 40 mL), and the combined ether extracts are washed repeatedly with water to remove any unreacted aminoalcohol. The ether containing the product is then dried over sodium sulfate, and the solvent is removed to leave the crude free ester. The free ester is converted to the hydrochloride salt by treatment with HCl in ether. The resulting solid is washed with ether and recrystallised from hot acetate/methanol to yield the title compound.

25

30

35

EXAMPLE 27

5 (±)-*trans*-[2-(4-morpholinyl)cyclohexyl]diphenylacetate
monohydrochloride
(Compound #27)

10 A solution of diphenylacetylchloride (2.95 g, 12.8 mmol) in chloroform (10 mL)
is slowly added to a solution of (±)-*trans*-[2-(4-morpholinyl)]cyclohexanol
prepared as in Example 4 (i) (2.25 g, 12.2 mmol) in chloroform (10 mL) under
nitrogen. The mixture is refluxed for 14 hours, and the crude ester is obtained
using the work-up procedure described in Example 26. It is dissolved in ether
15 (40 mL) and dichloromethane (5 mL) and treated with HCl in ether to precipitate
the salt, which is washed with ether and recrystallised from ethyl
acetate/methanol to yield the title compound.

EXAMPLE 28

20 (±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-3-trifluoromethylbenzoate
monohydrochloride
(Compound #28)

25 3-(Trifluoromethyl)benzoylchloride (2.96 g, 14.2 mmol), is dissolved in
chloroform (15 mL) and is refluxed for 60 hours with a solution of (±)-*trans*-[2-
(4-morpholinyl)]cyclohexanol (2.5 g, 13.5 mmol) in chloroform (12 mL) under
nitrogen. The solvent is removed *in vacuo*, and the residue is treated as in
Example 26 to give the crude free ester. This is purified by column
30 chromatography using silica gel and chloroform/ethyl acetate (9:1) as eluent.
The free ester is converted to the salt by treatment with HCl in ether. The
resulting solid is washed with ether and recrystallised from hot ethyl
acetate/methanol to yield the title compound.

EXAMPLE 29

[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]naphthyl-1-acetate
monohydrochloride
(Compound #29)

(i) 7-(Pyrrolidinyl)-1-oxaspiro[4.5]decan-8-ol is prepared by the method described in Example I(i), refluxing the corresponding epoxide (5 g, 32 mmol, prepared according to the method described in U.S. Patent No. 4,737,493) with pyrrolidine (2.28 g, 32 mmol) and water (5 mL) under nitrogen until GC or tlc analysis show the reaction to be complete. The produce is purified by full vacuum distillation.

(ii) The title compound is prepared according to the method described in Example 8 using 1-naphthylacetic acid (2 g, 10.8 mmol) and 7-(pyrrolidinyl)-1-oxaspiro[4.5]decan-8-ol (2.25 g, 10 mmol). The solvent is removed *in vacuo*, and the residue is treated as in Example 8 to leave the crude free ester. This is dissolved in ether (40 mL) and dichloromethane (5 mL) and treated with HCl in ether to precipitate the salt. It is washed with ether and recrystallised from ethyl acetate/methanol to yield the title compound.

Table 1

Compound	Formula	Calculated	Found
#1	C ₁₈ H ₂₄ NO ₃ Cl ₃	C 52.89, H 5.92, N 3.43%	C 52.59, H 5.78, N 3.37%
#3	C ₂₂ H ₃₀ NO ₃ Cl ₃	C 57.09, H 6.53, N 3.03%	C 57.05, H 6.31, N 3.08%
#4	C ₁₈ H ₂₄ NO ₄ Cl ₃	C 50.90, H 5.70, N 3.30%	C 50.88, H 5.70, N 3.25%
#5	C ₂₀ H ₂₈ NO ₃ Cl ₃	C 54.99, H 6.46, N 3.21%	C 54.31, H 6.49, N 3.26%
#6	C ₂₀ H ₂₆ NO ₃ Cl ₃	C 60.67, H 6.62, N 3.54%	C 60.69, H 6.57, N 3.58%
#7	C ₁₈ H ₃₂ NO ₂ Cl	C 65.53, H 9.78, N 4.25%	C 65.78, H 9.84, N 4.42%
#8	C ₂₂ H ₂₈ NO ₃ Cl	C 67.77, H 7.24, N 3.59%	C 67.47, H 7.21, N 3.57%
#9	C ₂₂ H ₂₈ NO ₃ Cl	C 67.77, H 7.24, N 3.59%	C 67.60, H 7.25, N 3.66%
#10	C ₁₈ H ₂₆ NO ₃ Cl	C 63.61, H 7.71, N 4.12%	C 63.50, H 7.56, N 4.17%
#11	C ₁₈ H ₂₄ NO ₃ Cl ₃	C 52.89, H 5.92, N 3.43%	C 52.75, H 5.90, N 3.40%
#12	C ₁₈ H ₂₅ N ₂ O ₅ Cl	C 56.18, H 6.55, N 7.28%	C 55.85, H 6.48, N 7.20%
#13	C ₂₃ H ₃₂ N ₂ O ₂ Cl ₂	C 62.87, H 7.34, N 6.38%	C 62.65, H 7.22, N 6.28%

	#14	C ₂₃ H ₃₁ N ₂ O ₂ Cl	C 68.56, H 7.75, N 6.95%	C 67.63, H 7.52, N 6.91%
	#15	C ₂₁ H ₂₆ NO ₃ Cl	C 67.10, H 6.97, N 3.73%	C 67.17, H 6.90, N 3.78%
	#16	C ₂₀ H ₂₆ NO ₃ ClS	C 60.67, H 6.62, N 3.54%	C 60.40, H 6.53, N 3.51%
	#17	C ₁₈ H ₂₅ NO ₃ ClBr	C 51.63, H 6.02, N 3.34%	C 51.56, H 6.01, N 3.34%
5	#18	C ₁₉ H ₂₄ NO ₃ Cl ₃	C 54.24, H 5.75, N 3.33%	C 54.33, H 5.78, N 3.32%

EXAMPLE 30

10

Assessment of Antiarrhythmic Efficacy

Antiarrhythmic efficacy was assessed by investigating the effect of a compound on the incidence of cardiac arrhythmias in conscious rats subject to coronary artery occlusion. Rats weighing 200-300 gms were subjected to preparative surgery and assigned to groups in a random block design. In each case, the animal was anesthetized with halothane during surgical preparation. The left femoral artery was cannulated for measurement of mean arterial blood pressure and withdrawal of blood samples. The left femoral vein was also cannulated for injection of drugs. The thoracic cavity was opened and a polyethylene occluder loosely placed around the left anterior descending coronary artery. The thoracic cavity was then closed. ECG was recorded by insertion of electrodes placed along the anatomical axis of the heart. All cannulae and electrode leads were exteriorized in the mid scapular region. In random and double-blind manner, about 0.5 to 2 hours post-surgery, an infusion of vehicle, or the compound to be tested was given. After 15 minutes infusion, the occluder was pulled so as to produce coronary artery occlusion. ECG, arrhythmias, blood pressure, heart rate and mortality were monitored for 30 minutes after occlusion. Arrhythmias were recorded as ventricular tachycardia (VT) and ventricular fibrillation (VF) and scored according to Curtis, M.J. and Walker, M.J.A., *Cardiovasc. Res.* 22:656 (1988) (see Table 2).

Table 2

	Score	Description
35	0	0-49 VPBs
	1	50-499 VPBs
	2	> 499 VPBs and/or 1 episode of spontaneously reverting VT or VF

- 3 > 1 episode of VT or VF or both (< 60s total combined duration)
 4 VT or VF or both (60-119s total combined duration)
 5 VT or VF or both (> 119s total combined duration)
 6 fatal VF starting at > 15 min after occlusion
 5 7 fatal VF starting at between 4 min and 14 min 59s after occlusion
 8 fatal VF starting at between 1 min and 3 min 59s after occlusion
 9 fatal VF starting < 1 min after occlusion

Where:

- 10 VPB = ventricular premature beats
 VT = ventricular tachycardia
 VF = ventricular fibrillation

15 Rats were excluded from the study if they did not exhibit pre-occlusion serum potassium concentrations within the range of 2.9-3.9 mM. Occlusion is associated with increases in R-wave height and "S-T" segment elevation; and an occluded zone (measured after death by cardiogreen dye perfusion) in the range of 25%-50% of total left-ventricular weight.

20 Table 3 describes the result of tests of the compounds described therein as values of arrhythmia score at a given dose in micromoles/kg/min.

Table 3

<u>Compound</u>	<u>Dose</u>	<u>AS</u>
Vehicle		
#2	20	7
#4	5	4
#5	4	3.6
#6	5	3
#8	2	1.8
#9	4	3
#10	10	1.3
#12	8	2.2
#15	12	3
#16	8	2
		0.3

EXAMPLE 31

Measurement of ECG Parameters

5 Rats weighing 200-250 gms were used in this example. Animals were anesthetized with 60 mg/kg pentobarbitone i.p. The carotid artery and jugular vein were cannulated for measurement of blood pressure and drug injection, respectively. ECG was recorded by insertion of electrodes placed
10 along the anatomical axis of the heart. All compounds were given as bolus injections.

Various ECG parameters were measured. The most sensitive of these for measuring sodium channel blockade is RSh (*J. Pharmacology Methods* 27:51-58 (1992)). Table 4 describes the results of the tests as ED₂₅
15 (micromoles/kg) which are the doses required to produce a 25% increase in the parameter measured (ne = not estimated). The increases in P-R interval, QRS interval and RSh indicate cardiac sodium channel blockade while the increase in Q-T interval indicates ancillary cardiac potassium channel blockade which is the
20 property of a type 1a antiarrhythmic.

Table 4

<u>Compound</u>	<u>RSh</u>	<u>PR</u>	<u>QRS</u>	<u>QT</u>
#1	10	17	ne	78
#2	11	23	ne	53
#3	140	140	130	160
#4	10	44	ne	ne
#5	36	43	66	45
#6	100	ne	ne	79
#7	6	84	ne	41
#8	11	20	42	57
#9	13	ne	ne	20
#10	16	ne	ne	25
#12	17	64	82	44
#15	80	340	90	ne
#16	23	230	690	33

EXAMPLE 32

Assessment of Sodium Channel Blockade

Rats were prepared according to the preceding procedure. Two silver stimulating electrodes were inserted through the chest wall and implanted in the left ventricle. Square wave stimulation was used to determine threshold current for capture, ventricular fibrillation threshold current, and effective refractory period (Howard, P.G. and Walker, M.J.A., *Proc. West. Pharmacol. Soc.* 33:123-127 (1990)). Table 5 contains ED₂₅ values for these indices of cardiac sodium channel blockade, where the ED₂₅ is the infusion rate in micromoles/kg/minute of compound required to elicit a 25% increase from control. The increases in refractoriness indicate ancillary blockade of potassium channels. The threshold current for capture is represented by "iT". The fibrillation threshold current is represented by "VFT". The effective refracting period is represented by "ERP".

Table 5

<u>Compound</u>	<u>iT</u>	<u>VFT</u>	<u>ERP</u>
#1	6	3	4
#2	5	2	3
#5	5	4	4
#6	12	9	7
#8	5	2	4
#9	34	7	8
#10	15	11	8
#12	4	5	6
#15	33	18	20
#16	15	2	5

EXAMPLE 33

Voltage Clamp Tests

5 In order to directly measure cardiac sodium channel blockade, standard whole cell voltage clamp tests were carried out using single isolated adult rat cardiac cells. Results in the form of EC₅₀ values, which are micromolar concentrations required to produce a 50% reduction in sodium
10 current, are presented in Table 6.

Table 6Compound

#1	200
#2	50
#3	>150
#4	>500
#6	150
#8	80
#9	>100
#10	>300
#12	>300
#15	ne
#16	225

15

EXAMPLE 34

Nerve Conduction Blockade

20

In order to establish the potency of the compounds to block nerve conduction, two *in vitro* assays were used. The first was the phrenic nerve diaphragm of the rat. The second was the hypogastric nerve vas deferens. Nerve blockade in these preparations is indicative of local anesthetic activity. Compounds were administered to the bath solution in increasing concentrations

while the preparations were being stimulated with regular electrical impulses. Table 7 shows the micromolar concentrations at which the twitch response from electrical stimulation is reduced to 50% of control value.

Table 7

	Compound				
	#1	#2	#3	#5	#6
Phrenic nerve diaphragm	90	120	200	180	30
Hypogastric nerve vas deferens	40	25	300	100	100

EXAMPLE 35


Opioid-like Effects

During *in vivo* tests of compound #6, it was observed that this compound induced opioid-like effects on the central nervous system of the rat. These effects were completely blocked by naloxone, an inhibitor of mu, delta and kappa opioid agonists. This provides evidence that some of the compounds may have analgesic activity through activation of opioid receptors. Examination of members of this series of compounds by analgesic assays and binding studies show the presence of opioid binding and activity.

From the foregoing, it will be evident that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention.

Claims

1. An enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:




(I)

wherein X is a direct bond;
 or $-(CH_2)_n-Y-$, where $n = 1, 2$, or 3 , and Y is a direct bond, O or S;
 or $-CH(R_{12})-Y-$, where R_{12} is alkyl of from one to six carbon atoms, a saturated carbocyclic ring of from three to six carbon atoms, phenyl or benzyl, and Y is a direct bond, O or S;
 or $-C(R_{13})=CH-$, where R_{13} is hydrogen, alkyl of from one to six carbon atoms, or phenyl;

R_1 and R_2 are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

or R_1 and R_2 , when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II:



(II)

where m is an integer from three to eight, and said ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of


one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and said heterocyclic ring containing not more than two nitrogen ring atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

R₃ and R₄ are independently attached to the cyclohexane ring at the 3-, 4-, 5-, or 6-positions, and are independently hydrogen, hydroxy, alkyl of one to six carbon atoms or alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; and


A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from formulae III, IV, V, VI, VII or VIII:



(III)


where R₅, R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl, and when X is a direct bond at least one of R₅, R₆ and R₇ is a hydroxy, fluorine, chlorine, bromine, trifluoromethyl, alkyl of from one to six carbon atoms, or aryl substituent, and when X is -CH=CH-, and R₁ and R₂ when taken together with the nitrogen atom to which they are attached, form a N-phenylpiperazine ring, and R₃ and R₄ are hydrogen, at least one of R₅, R₆ and R₇ is a substituent other than hydrogen;

66




(IV)

where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;




(V)

where R₈ and R₉ are defined as above;




(VI)

where R₁₀ is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms;



(VII)

only when X is a direct bond;




(VIII)

only when X is a direct bond;

with the proviso that, when X is $-(CH_2)_n-Y-$, and $n = 1$, and Y is a direct bond, and R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may not be 4-thianaphthenyl.

2. An enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



wherein X is a direct bond;

or $-(CH_2)_n-Y-$, where $n = 1$ and Y is a direct bond, O or S;

or $-CH(R_{12})-$, where R_{12} is alkyl of from one to six carbon atoms;

or $-C(R_{13})=CH-$, where R_{13} is hydrogen;


R_1 and R_2 are defined as in claim 1;

R_3 and R_4 are defined as in claim 1; and

A is defined as in claim 1;

with the proviso that, when X is $-(CH_2)_n-Y-$, and $n = 1$, and Y is a direct bond, and R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may not be 4-thianaphthenyl.

3. An enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



wherein X is $-(CH_2)_n-Y-$, where $n = 1$ and Y is a direct bond or O;
or $-CH(R_{12})-$, where R_{12} is alkyl of from one to six carbon atoms;

R_1 and R_2 are defined as in claim 1;

R_3 and R_4 are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom; and

A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from:

formula III where R_5 , R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl;

or formula IV where R_8 and R_9 are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;


or formula V where R_8 and R_9 are defined as above;

or formula VI where R_{10} is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z

is CH_2 , O, S, or N-R_{11} where R_{11} is hydrogen or alkyl of one to six carbon atoms;

with the proviso that, when X is $-(\text{CH}_2)_n\text{-Y-}$, and $n = 1$, and Y is a direct bond, and R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may not be 4-thianaphthenyl.

4. An enantiomer or geometric isomer of a compound of formula IX, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



(IX)

wherein R_1 and R_2 , when taken together with the nitrogen atom to which they are attached,

form a ring according to formula II, where m is an integer from three to eight, and said ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and said heterocyclic ring containing not more than two nitrogen ring forming atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5- positions, and are independently hydrogen, methoxy, or are points of attachment of a five-membered oxaspiran ring; and

A is a saturated carbocyclic ring of three to six carbon atoms, or is selected from:


formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R₆ and R₇ is a substituent other than hydrogen;

or formula IV where R₈ and R₉ are hydrogen;

or formula V where R₈ and R₉ are hydrogen;

or formula VI where R₁₀ is hydrogen, and Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or methyl.

5. An enantiomer or geometric isomer of a compound of formula X, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



(X)

wherein R₁ and R₂ are defined as in claim 4;

R₃ and R₄ are defined as in claim 4; and

A is a saturated carbocyclic ring of from three to six carbon atoms, or is selected from:

formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R_6 and R_7 is a substituent other than hydrogen;


or formula IV where R_8 and R_9 are hydrogen;

or formula V where R_8 and R_9 are hydrogen;

or formula VI where R_{10} is hydrogen, and Z is CH_2 , O, S, or N- R_{11} where R_{11} is hydrogen or methyl;

with the proviso that when R_1 and R_2 , taken together with the nitrogen atom to which they are attached, form a pyrrolidinyl ring, and R_3 and R_4 are hydrogen, A may not be 4-thianaphthenyl.

6. An enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



(I)

wherein X is a direct bond or $-CH=CH-$;

R_1 and R_2 are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

or R₁ and R₂, when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II where m is an integer from three to eight;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5-positions, and are independently hydrogen, methoxy, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom; and

A is a saturated carbocyclic ring of from three to six carbon atoms, or is selected from:

formula III where R₅ is hydrogen, and R₆ and R₇ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, trifluoromethyl, methyl or ethyl, and at least one of R₆ and R₇ is a substituent other than hydrogen;

or formula IV where R₈ and R₉ are hydrogen;

or formula V where R₈ and R₉ are hydrogen;


or formula VI where R₁₀ is hydrogen, and Z is O, S, or N-R₁₁ where R₁₁ is hydrogen or methyl;

or formula VII when X is a direct bond;

or formula VIII when X is a direct bond.

7. An enantiomer or geometric isomer of a compound of formula XI, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:

73



(XI)

wherein R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a ring according to formula II where m is an integer from three to eight; and

A is selected from:


formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R_6 and R_7 is a substituent other than hydrogen;

or formula IV where R_8 and R_9 are hydrogen;

or formula V where R_8 and R_9 are hydrogen;

or formula VI where R_{10} is hydrogen, and Z is O or S.

8. An enantiomer or geometric isomer of a compound of formula XII, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



(XII)

wherein R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a ring according to formula II where m is an integer from three to eight; and

A is selected from:

formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy, and at least one of R_6 and R_7 is a substituent other than hydrogen;


or formula IV where R_8 and R_9 are hydrogen;

or formula V where R_8 and R_9 are hydrogen;

or formula VI where R_{10} is hydrogen, and Z is O or S;

with the proviso that, when R_1 and R_2 taken together with the nitrogen atom to which they are attached form a pyrrolidinyl ring, A may not be 4-thianaphthenyl.

9. An enantiomer or geometric isomer of a compound of formula XIII, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:




(XIII)

wherein X is a direct bond, *trans*-CH=CH-, -CH₂- or -CH₂-O-;

R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, complete a ring selected from pyrrolidiny, piperidiny, hexahydroazepiny, morpholiny, methylpiperaziny or 3-azabicyclo[3.2.2]nonany; and

A is selected from cyclohexyl, 3,4-dichlorophenyl, 4-bromophenyl, 1-naphthyl, 2-naphthyl or 3-thianaphthenyl.

10. An enantiomer or geometric isomer of a compound of formula XIV, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



(XIV)

wherein X is a direct bond;

or $-(CH_2)_n-Y-$, where $n = 1, 2$, or 3 , and Y is a direct bond, O or S;


or $-CH(R_{12})-Y-$, where R_{12} is alkyl of from one to six carbon atoms, a saturated carbocyclic ring of from three to six carbon atoms, phenyl or benzyl, and Y is a direct bond, O or S;

or $-C(R_{13})=CH-$, where R_{13} is hydrogen, alkyl of from one to six carbon atoms, or phenyl;

R_1 and R_2 are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

or R_1 and R_2 , when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II:



where m is an integer from three to eight, and said ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and said heterocyclic ring containing not more than two nitrogen ring atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;


R_3 and R_4 are independently attached to the cyclohexane ring at the 3-, 4-, 5-, or 6-positions, and are independently hydrogen, hydroxy, alkyl of one to six carbon atoms or alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; and

A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from formulae III, IV, V, VI, VII or VIII:




where R_5 , R_6 and R_7 are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl, and when X is a direct bond at least one of R_5 , R_6 and R_7 is a substituent other than hydrogen;

77




(IV)

where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;




(V)

where R₈ and R₉ are defined as above;



(VI)


where R₁₀ is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms;



(VII)

only when X is a direct bond;


78



(VIII)

only when X is a direct bond.

11. An enantiomer or geometric isomer of a compound of formula XIV, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



(XIV)

wherein X is a direct bond;

or $-(CH_2)_n-Y-$, where $n = 1$, and Y is a direct bond, O or S;

or $-CH(R_{12})-$, where R_{12} is alkyl of from one to six carbon atoms;


or $-C(R_{13})=CH-$, where R_{13} is hydrogen;

R_1 and R_2 are defined as in claim 10;

R_3 and R_4 are independently attached to the cyclohexane ring at the 4- or 5- positions, and are independently hydrogen, alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen atom; and

A is defined as in claim 10.

12. An enantiomer or geometric isomer of a compound of formula XV, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



wherein X is a -CH₂- or -CH₂-O-;

R₁ and R₂, when taken together with the nitrogen atom to which they are attached, form a ring according to formula II, where m is an integer from three to eight, and said ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;

or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and said heterocyclic ring containing not more than two nitrogen ring forming atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

R₃ and R₄ are independently attached to the cyclohexane ring at the 4- or 5- positions, and are independently hydrogen, methoxy, or are points of attachment of a five-membered oxaspiran ring; and

A is a saturated carbocyclic ring of from three to six carbon atoms, or is selected from:


formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy;

or formula IV where R_8 and R_9 are hydrogen;

or formula V where R_8 and R_9 are hydrogen;

or formula VI where R_{10} is hydrogen, and Z is CH_2 , O, S, or N- R_{11} where R_{11} is hydrogen or methyl.

13. An enantiomer or geometric isomer of a compound of formula XVI, or a solvate or pharmaceutically acceptable salt thereof, said compound of the formula:



(XVI)

wherein X is a $-CH_2-$ or $-CH_2-O-$;

R_1 and R_2 , when taken together with the nitrogen atom to which they are attached, form a ring according to formula II where m is an integer from three to eight; and

A is selected from:

formula III where R_5 is hydrogen, and R_6 and R_7 are independently hydrogen, hydroxy, fluorine, chlorine, bromine, nitro, trifluoromethyl, methyl, ethyl, methoxy, or ethoxy;

or formula IV where R_8 and R_9 are hydrogen;

or formula V where R₈ and R₉ are hydrogen;

or formula VI where R₁₀ is hydrogen, and Z is O or S.

14. A compound selected from the group consisting of:

(±)-*trans*-[2-(1-pyrrolidinyl)cyclohexyl](3,4-dichlorophenoxy)acetate;
(±)-*trans*-[2-(1-pyrrolidinyl)cyclohexyl](1-naphthoxy)acetate;
(±)-*trans*-{2-[N-(3-azabicyclo[3.2.2]nonyl)]cyclohexyl}(3,4-dichlorophenoxy)acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl](3,4-dichlorophenoxy)acetate;
(±)-*trans*-[2-(1-hexahydroazepinyl)cyclohexyl](3,4-dichlorophenoxy)acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]benzo[b]thiophene-3-acetate;
(±)-*trans*-[2-(1-pyrrolidinyl)cyclohexyl](cyclohexyl)acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]naphthyl-1-acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]naphthyl-2-acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]phenylacetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl](3,4-dichlorophenyl)acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl](4-nitrophenyl)acetate;
(±)-*trans*-[2-(4-methyl-1-piperazinyl)cyclohexyl]naphthyl-2-acetate;
(±)-*trans*-[2-(4-methyl-1-piperazinyl)cyclohexyl]naphthyl-1-acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-1-naphthoate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]benzo[b]thiophene-4-acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl](4-bromophenyl)acetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-3,4-dichlorocinnamate;
(±)-*trans*-[2-(1-piperidinyl)cyclohexyl]thiophenoxyacetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-2-(1-naphthyl)propionate;
(±)-(1 α ,2 β ,4 β ,5 β)-[4,5-dimethoxy-2-(1-pyrrolidinyl)cyclohexyl]benzofuran-2-acetate;
(±)-*trans*-{2-[bis(2-methoxyethyl)amino]cyclohexyl}fluorene-9-carboxylate;
S-{(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]}naphthyl-1-thioacetate;
S-{(±)-*trans*-{2-[bis(2-methoxyethyl)amino]cyclohexyl}}-4-bromothiobenzoate;
(±)-*trans*-[2-(diisopropylamino)cyclohexyl]acenaphthene-1-carboxylate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]diphenylacetate;
(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]-3-trifluoromethylbenzoate;
[7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl]naphthyl-1-acetate;
S-{(±)-*trans*-[2-(4-morpholinyl)cyclohexyl]}(3,4-dimethoxyphenyl)thioacetate; and
pharmaceutically acceptable salts thereof.

15. A composition comprising a compound according to any one of claims 1-14 in combination with a pharmaceutically acceptable carrier or diluent.

16. A compound according to any one of claims 1-9 and 14 for use in a method for treating arrhythmia in a warm-blooded animal.


17. A compound according to any one of claims 1-9 and 14 for use in a method for blocking ion channels in a warm-blooded animal.

18. A compound according to any one of claims 10-13 for use in a method for treating arrhythmia in a warm-blooded animal.

19. A compound according to any one of claims 10-13 for use in a method for blocking ion channels in a warm-blooded animal.

20. A compound according to any one of claims 1-14 for use in a method for blocking ion channels *in vitro*.

21. Use of an enantiomer or geometric isomer of a compound of formula I, or a solvate or pharmaceutically acceptable salt thereof, in the manufacture of a medicament to treat arrhythmia, said compound of the formula:



wherein X is a direct bond;

or $-(CH_2)_n-Y-$, where $n = 1, 2, \text{ or } 3$, and Y is a direct bond, O or S;


or $-CH(R_{12})-Y-$, where R_{12} is alkyl of from one to six carbon atoms, a saturated carbocyclic ring of from three to six carbon atoms, phenyl or benzyl, and Y is a direct bond, O or S;

or $-C(R_{13})=CH-$, where R_{13} is hydrogen, alkyl of from one to six carbon atoms, or phenyl;

R₁ and R₂ are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

or R₁ and R₂, when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II:



where m is an integer from three to eight, and said ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;


or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and said heterocyclic ring containing not more than two nitrogen ring atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

R₃ and R₄ are independently attached to the cyclohexane ring at the 3-, 4-, 5-, or 6-positions, and are independently hydrogen, hydroxy, alkyl of one to six carbon atoms or alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; and


A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from formulae III, IV, V, VI, VII or VIII:

84




(III)

where R₅, R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl;




(IV)

where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;



(V)


where R₈ and R₉ are defined as above;



(VI)


where R₁₀ is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms;

85



(VII)


only when X is a direct bond;



(VIII)

only when X is a direct bond.

22. Use of an enantiomer or geometric isomer of a compound of formula XIV, or a solvate or pharmaceutically acceptable salt thereof, in the manufacture of a medicament to treat arrhythmia, said compound of the formula:



(XIV)

wherein X is a direct bond;

or $-(CH_2)_n-Y-$, where $n = 1, 2$, or 3 , and Y is a direct bond, O or S;


or $-CH(R_{12})-Y-$, where R_{12} is alkyl of from one to six carbon atoms, a saturated carbocyclic ring of from three to six carbon atoms, phenyl or benzyl, and Y is a direct bond, O or S;

or $-C(R_{13})=CH-$, where R_{13} is hydrogen, alkyl of from one to six carbon atoms, or phenyl;

R₁ and R₂ are independently hydrogen, alkyl of three to eight carbon atoms, alkoxyalkyl of three to eight carbon atoms, or aralkyl of seven to twelve carbon atoms;

or R₁ and R₂, when taken together with the nitrogen atom to which they are attached,

form a ring denoted by formula II:



where m is an integer from three to eight, and said ring may be substituted at any one carbon atom by hydroxy, oxo, alkyl of one to three carbon atoms or alkoxy of one to three carbon atoms, or may be fused at two adjacent carbon atoms with an aromatic or aliphatic carbocyclic ring of six carbon atoms;


or complete a saturated monocyclic nitrogen heterocyclic ring of five to eight ring atoms, containing only carbon, nitrogen and optionally oxygen ring atoms, and said heterocyclic ring containing not more than two nitrogen ring atoms, the second nitrogen being optionally substituted with an alkyl group of one to six carbon atoms or a phenyl ring;

or complete a ring selected from 3-azabicyclo[3.2.2]nonan-3-yl, 2-azabicyclo[2.2.2]octan-2-yl, 3-azabicyclo[3.1.0]hexan-3-yl, or 3-azabicyclo[3.2.0]heptan-3-yl;

R₃ and R₄ are independently attached to the cyclohexane ring at the 3-, 4-, 5-, or 6-positions, and are independently hydrogen, hydroxy, alkyl of one to six carbon atoms or alkoxy of one to six carbon atoms, or are points of attachment of a spiro five- or six-membered heterocyclic ring containing one oxygen or sulfur atom; and


A is an alkyl group of five to twelve carbon atoms, or is a saturated carbocyclic ring of three to six carbon atoms, or is selected from formulae III, IV, V, VI, VII or VIII:

87




(III)

where R₅, R₆ and R₇ are independently hydrogen, hydroxy, amino, fluorine, chlorine, bromine, nitro, trifluoromethyl, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms, or aryl;




(IV)

where R₈ and R₉ are independently hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of one to six carbon atoms, alkoxy of one to six carbon atoms or aryl;



(V)


where R₈ and R₉ are defined as above;



(VI)


where R₁₀ is hydrogen, hydroxy, fluorine, chlorine, bromine, alkyl of from one to six carbon atoms, alkoxy of from one to six carbon atoms, or aryl; Z is CH₂, O, S, or N-R₁₁ where R₁₁ is hydrogen or alkyl of one to six carbon atoms;

88



(VII)

only when X is a direct bond;



(VIII)

only when X is a direct bond.

INTERNATIONAL SEARCH REPORT

Internat. Application No
PCT/CA 94/00513

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D295/08 C07D223/14 C07D333/56 C07D333/54 C07D307/80
C07C219/24 C07C323/30 C07D307/94 A61K31/215

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US,A,2 954 380 (SEYMOUR L. SHAPIRO ET AL) 27 September 1960 *Complete document, particularly column 1 : lines 15-34*	1-22
P,A	FR,A,2 689 893 (RHONE-POULENC RORER) 15 October 1993 *Page 15-17: claims*	1,21,22



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

12 January 1995

Date of mailing of the international search report

19. 01. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Luyten, H

4672

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 94/00513

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-2954380		NONE	
FR-A-2689893	15-10-93	NONE	